

Open-Source Value Project (OSVP) IVI RA-Model Version 1.0

Review Overview and Plan for Revision

September 20, 2018



This work is licensed under the Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



About the Innovation and Value Initiative

The Innovation and Value Initiative (IVI) is a multi-stakeholder initiative that seeks to improve the way value is measured and rewarded in the healthcare system to promote the development and use of high value interventions that advance human health. To achieve this mission, IVI pursues the following goals:

- Establish best practices for measuring the real-world value of healthcare technologies using both existing and innovative scientific methods;
- Provide a range of marketplace stakeholders – including patients, consumers, providers, healthcare systems, and payers – with salient, accurate, and actionable information about value in healthcare;
- Develop and test innovative approaches to link healthcare spending to value.

The IVI is hosted by Precision Health Economics, a health economics consultancy. IVI's direction and research agenda are determined in collaboration with its Strategic Advisory Panel, which includes representatives from patient advocacy organizations, pharmaceutical firms, academia, insurers, and health systems. All funding supports IVI's overall activities, with no funding or funder tied to specific activities or research projects.

Funding

This study was funded by the Innovation and Value Initiative, a non-profit collaboration among thought leaders in academia, patient advocacy organizations, payers, pharmaceutical companies, providers, delivery systems and other organizations dedicated to advancing the science and application of value assessment in health care.

Table of Contents

About the Innovation and Value Initiative.....	2
Funding	2
Table of Contents.....	3
Executive Summary.....	4
Introduction.....	5
Overview of Process for Updating the IVI-RA Model.....	5
Public Release of the IVI-RA Model.....	5
Solicitation of Public Comment.....	6
Technical Expert Panel (TEP) Review and Priority-Setting.....	7
Methods.....	7
Results.....	9
Final Recommendations of Technical Expert Panel	12
Anticipated Timeline for All Revisions	13
Appendix A. Public Comments on Initial Release of IVI-RA Modeling Platform.....	14
Appendix B. IVI-RA Model Technical Expert Panel Membership.....	15
Appendix C. Delphi Surveys.....	17
Appendix D. Delphi Survey Results.....	39
References	44



Executive Summary

Inspired by the open-source software development process, IVI's Open-Source Value Project (OSVP) uses a transparent, collaborative, and iterative process to develop flexible, open-source value models in specific diseases. These models – and the process of developing them – serve as a hands-on laboratory for advancing the science of value assessment, engaging stakeholders in constructive debate, and ultimately building consensus around how value is measured in a diverse and decentralized health care system. Each value model release is followed by: public feedback as part of an open comment period; review and discussion of the feedback by the Technical Expert Panel (TEP); and identification of specific recommendations for changes using a modified Delphi process. To ensure transparency, IVI documents, summarizes, and makes available the details of these deliberations and decisions.

IVI released its initial version of a disease-specific OSVP model focused on Rheumatoid Arthritis (the IVI-RA model) in November 2017, after which IVI invited public comment through February 16, 2018. Upon the conclusion of the public comment period, IVI engaged a third-party Technical Expert Panel (TEP) comprised of leaders in health economics, epidemiology, rheumatology, and patient communities to review the public comments and establish priorities for model improvement through a teleconference and a two-part modified Delphi survey.

Several priorities emerged from TEP deliberation that will shape IVI's approach to developing Version 2.0 of the IVI-RA model, including:

1. IVI will update both the systematic literature review and the network meta-analysis to ensure the model reflects the most recent clinical evidence including newly available treatments.
2. IVI will expand model flexibility by adding more utility functions, allowing for treatment specific discontinuation, potentially including additional outcomes correlated with HAQ based on literature, and adding biosimilars as an option;
3. IVI will seek out opportunities to use real-world data sources;

IVI will release an updated Version 2.0 of the IVI-RA model in the second quarter of 2019.



Introduction

With the Open-Source Value Project (OSVP), IVI strives to maximize both the relevance and credibility of value assessment in the context of the United States' decentralized decision-making environment by developing and providing access to flexible open-source decision models for value assessment. These interactive models have two primary objectives: (i) to enable a more constructive dialogue regarding value assessment between stakeholders (e.g. patients, payers, providers, and manufacturers) with different beliefs about relevant clinical data, modeling approaches, and value perspectives; and (ii) to provide local decision-makers with credible value assessments that reflect the local setting and is based on the latest evidence while accounting for all scientific uncertainty (due to patient heterogeneity, gaps in evidence, and different modeling beliefs). OSVP facilitates iterative development and collaboration between multiple clinical and methodological experts, and evolves as new clinical evidence and scientific insights become available.

IVI's first disease-specific OSVP model in Rheumatoid Arthritis (RA), the IVI-RA model, is a discrete-time individual patient simulation that simulates outcomes for individual patients. The model measures clinical outcomes (e.g., change in functional status over time, rate of adverse events), costs and other health economics measures (e.g., quality-adjusted life years), and other factors in order to measure a treatment's value. Value is measured both using a more traditional cost-effectiveness analysis (CEA) as well as a multi-criterion decision analysis (MCDA) framework which allows differences in patient preferences to influence the preferred treatment.

Overview of Process for Updating the IVI-RA Model

IVI's Open Source Value Project (OSVP) aims to build an open, collaborative, and consensus-based process for the development of tools for value assessment. Models developed by the OSVP process are iterative, evolving as the science of value assessment advances, and as new evidence becomes available.

The public release and update of OSVP models follow a four-step process:

1. Release of the model into the public domain
2. Solicitation of feedback through a public comment period
3. Technical Expert Panel (TEP) review of public feedback and prioritization of recommended updates using 2-part Delphi approach
4. Implementation of recommended changes by IVI

The OSVP process ensures that ongoing model development is driven by the feedback and suggestions of diverse stakeholders engaged in the public comment process, with the objective guidance of the TEP. Each of these steps, the resulting recommendations of the TEP, and the planned updates to the IVI-RA model are described below.

Public Release of the IVI-RA Model

In November of 2017, IVI released its first OSVP model in RA. The model examines outcomes and the relative value of drug therapies for moderate to severe RA. Version 1.0 of the RA model reflects a range of scientifically defensible approaches (i.e., 384 possible model structures), is highly customizable, includes multiple novel approaches to value assessment (i.e., MCDA), and additionally incorporates 'innovative' sources of value. A more in-depth description of the IVI-RA model methodology is [available online](#). [1]

Solicitation of Public Comment

The public comment period began in November of 2017 following the release of the model, and concluded in mid-February of 2018. Substantive comments were received from a diverse range of stakeholders, which included patient advocacy organizations, academics, physicians, non-profits, and life-science companies.

Feedback was received from the following organizations: AbbVie, Arthritis Foundation, Bristol-Myers Squibb, Genentech, ICER, Institute for Patient Access, Janssen, Pfizer, American College of Rheumatology Rheum-PAC, Sanofi-Genzyme and Regeneron, and Tufts University (see Appendix A for complete text). These comments generally focused on the following topics: model outcomes, model structure, evidence considered, treatment efficacy, and value tool. Primary takeaways regarding each of these categories are listed below:

Model outcomes:

- IVI should incorporate additional outcomes such as cardiovascular outcomes, caregiver burden, nursing home stays, additional patient reported outcomes (PROs)

Model structure:

- HAQ rebound: After a treatment failure, the HAQ should rebound by increasing an amount *equal* to its initial 6-month decrease (the model currently assumed a rebound ranging from 70% to 100%)
- DAS28: IVI should model the progression of DAS28 over time as well as HAQ
- Modified Total Sharp Score (mTSS): Although it is expected to be modest, IVI should model the *independent* effect of the mTSS on HAQ, since it adds clinical validity to the model
- Mortality: Association between HAQ and mortality should be based solely on patients with radiographic data (rather than all patients) from [Wolfe et al. \(2003\)](#); [2] it is unclear whether progression of HAQ (rather than just baseline HAQ) impacts mortality

Evidence considered:

- Including real-world data:
 - IVI should develop a partnership with a developer of an established observational dataset
 - The patient population should be based on registry data, rather than a summary of data from the literature
 - Patient preferences for treatments should be based on actual patient outcomes
- Drug prices:
 - WAC is inappropriate because it does not represent actual transaction prices or include prompt pay or other discounts, rebates, or reductions in price (note that the IVI-RA model does include discounts by default)
 - National Average Drug Acquisition Cost (NADAC) is a more appropriate measure of drug price than the WAC

Treatment efficacy:

- Rate of HAQ increase is uncertain beyond length of follow-up data (e.g., 15 years); a constant HAQ degradation rate should therefore be used after 15 years
- Unclear why treatment efficacy assumed to decline after *first* previous treatment failures by a constant factor; estimates should be based on available RCTs or observational data sources instead
- Serious infection rates should not be assumed equal across tDMARDs as there is real-world data that suggests otherwise (RCT evidence is less clear)
- The analysis should allow for subgroup analyses – e.g., patients with poor prognostic factors or more severe disease

Value tool:

- Default sequences: It is unclear how the default sequence is chosen, and whether it is the most relevant sequence to compare
- Change in functional status: It is difficult to interpret the HAQ disability index figure
- MCDA: The slider scales on the MCDA page are not intuitive; clarification is needed on how outcomes were placed on a common scale and the rationale for default settings
- Value of a QALY: The default willingness to pay per QALY of \$150,000 seems arbitrary and very high; results may therefore be misleading

Technical Expert Panel (TEP) Review and Priority-Setting

Methods

Upon the conclusion of the public comment period, IVI engaged a third-party Technical Expert Panel (TEP) comprised of five leaders in health economics, epidemiology, rheumatology, and patient communities to review all public comments and establish priorities for revision for the model's next iteration. See Appendix B for more information about the members of the TEP.

IVI engaged with the TEP across two phases between mid-February and end of May. In the first phase, the TEP conducted a 1-hour teleconference discussion to overview and qualitatively synthesize all public comments received in mid-February. In the second phase, members of the TEP participated in a modified Delphi process aimed at achieving consensus around finalizing priorities for revision. Consisting of two sequential surveys designed to identify areas of greater/lesser consensus and to measure priority of various recommendations, this approach was intended to provide objective and quantitative priorities for revisions.

The two-part modified Delphi process was structured as follows:

1. Administration of Survey #1:

- a. Each suggested change derived from the public comments and TEP recommendations at the teleconference was categorized by domain (model outcomes, model structure, treatment efficacy, evidence considered, value tool), and listed in a survey
- b. TEP members were asked to complete the survey by rank ordering each suggested change according to priority (1-4, with 4 being the highest priority), and elaborating their reasoning for their respective selections (See Appendix C for the full survey).

- c. In addition, to providing a quantitative 1 to 4 priority ranking, TEP members were also asked to provide the rationale behind their ranking.

2. IVI Internal Analysis

- a. IVI team analyzed the results of the first survey internally to determine areas of top priority, and areas where agreement was not reached. Items that were identified as top priority (mean priority score >3.5) were based on the priority value across all TEP responses. Areas of disagreement were the ones where the standard deviation of responses across TEP members was highest ($SD >0.5$).

3. Administration of Survey #2

- a. TEP members were asked to complete a follow-up survey that summarized and informed respondents on the areas of agreement from Survey #1, and solicited further feedback for areas where agreement was not reached. Specifically, the TEP was asked to (1) confirm whether they agreed that the recommendations with the highest mean priority value reflected their opinions, and (2) re-rank the recommendations where the standard deviation of priorities was highest across TEP members.
- b. For each recommendation, reviewers were able to view the average priority score across TEP members in Survey #1 as well as the anonymized rationale for why all TEP members selected those priority levels.
- c. IVI then analyzed the results to identify highest-priority updates based on all responses from both Survey #1 and Survey #2. High priority responses from Survey #1 were included in the final list unless a majority of the TEP decided that this was no longer a priority. Among the recommendations where there was uncertainty in Survey #1, IVI required the mean priority score in Survey #2 to be at least 3.0.

Results

Delphi Survey #1

The top priorities and areas of disagreement are summarized in Tables 1 and 2. See Appendix D for a complete table of results from Delphi Survey #1.

Overall, the results from the first Delphi Survey indicated that incorporating additional treatment options (i.e., triple therapy, biosimilars), expanding model functionality (i.e., including additional utility functions and allowing for treatment-specific discontinuation), and utilizing real-world data sources (i.e., CORRONA) were top priority revisions for the next iteration of the RA model. The inclusion of additional adverse events (i.e., tuberculosis infection risk) and available IV and SC formulations for all biologic DMARDs were deemed to be lower priority ($<2.8/4.0$).

While most respondents suggested the importance of including additional outcomes in the model (i.e., fractures, fatigue, cardiovascular events, long-term nursing home stays, caregiver burden), the results of the survey conveyed that there was considerable variability in regards to which additional outcomes should ultimately be incorporated. The presentation of HAQ simulated pain scores as a separate outcome also garnered reasonable disagreement among respondents.

Delphi Survey #2

The final top priorities and residual areas of disagreement are summarized in Tables 3 and 4. See Appendix D for a complete table of results from Delphi Survey #2.

In regards to top priorities for revision, the results from the second Delphi Survey were largely consistent with the results yielded from the first survey. Respondents reiterated the importance of including additional therapy options, expanding model functionality, and investing in the acquisition of patient-level, real world data. Additionally, TEP members also emphasized the need to utilize the NADAC to estimate the cost of drugs.

After the solicitation of further feedback, the areas of disagreement from the administration of the second survey differed from what was seen in the first. The inclusion of both an additional model structure, DAS-28, as an alternative measure of disease progression to HAQ, and the option of continuing with the same tDMARD despite experiencing a serious infection emerged as areas of disagreement.

Table 1: Recommended model improvements ranked highest priority in Delphi Survey #1

Top priorities based off of Average Score	Avg. score Scale (1-4)	Rank
The model should provide an option to select biosimilars in addition to the currently available biologic DMARDs.	3.8	1
Triple therapy should be included as a treatment option in the model.	3.8	1
As an alternative to modeling time-to-treatment-discontinuation during the maintenance phase with a specific tDMARD (>6 months) conditional upon the level of response or disease activity at 6 months with the same tDMARD, the model should provide the option to determine treatment discontinuation conditional upon HAQ or disease activity level during the maintenance phase.	3.8	3
In addition to probabilistic sensitivity analysis available with the current version of the model, the model should also present tornado diagrams, or another way of presenting the contribution of different model components/parameters on the uncertainty in model outcomes and cost-effectiveness estimates.	3.8	3
Additional utility functions beyond those already incorporated in the model should be included.	3.7	5

Table 2: Areas of greatest TEP disagreement identified in Delphi Survey #1

Top areas of disagreement based off of Standard Deviation (SD)	SD	Rank
In the model pain scores are estimated based on HAQ scores given their association. These simulated pain scores need to be presented separately as they are an important outcome.	1.3	1
Cardiovascular events should be included in the model.	1.2	2
tDMARD specific serious infection rates need to be incorporated.	1.2	2
Fatigue should be added as an outcome in the model.	1.1	4
The HAQ score should rebound by 100% at treatment discontinuation.	1.1	5

Table 3: Recommended model improvements ranked highest priority in Delphi Survey #2

Top priorities based off of Average Score	Avg. score	Rank
Sarilumab should be included in the model.	3.8	1
Long-term discontinuation rates should be drug specific.	3.8	1
The model should continue to allow changes in HAQ over time to affect mortality.	3.8	1
The NADAC should be used instead of the WAC (along with adjustments for rebates and discounts) to estimate drug acquisition costs.	3.3	4
IVI should spend resources to acquire patient-level real-world data.	3.3	4

Table 4: Areas of greatest TEP disagreement identified in Delphi Survey #2

Top areas of disagreement based off of Standard Deviation (SD)	SD	Rank
Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ.	1.1	1
Fractures should be included in the model.	0.7	2
The model should provide the option to continue with the same tDMARD despite experiencing a serious infection.	0.7	2
Treatment effects among tDMARD experienced patients should be modeled as a function of the number of previous tDMARDs received.	0.7	2
The NADAC should be used instead of the WAC (along with adjustments for rebates and discounts) to estimate drug acquisition costs.	0.5	5

Final Recommendations of Technical Expert Panel

After participation in the teleconference and the completion of both Delphi surveys, the recommendations in Table 5 emerged as the most salient recommendations from the TEP, and will ultimately serve as IVI priorities for revision.

Note: TEP recommendations are based solely on scientific merit, and to determine and finalize priorities for revision, the IVI team considers priority rankings derived from TEP deliberation in addition to data availability, budget constraints and other factors to identify which items are feasible. However, the TEP recommendations serve as the most important criteria for updating the model.

Table 5: Final recommendations from TEP and IVI planned responses

TEP Recommendation	IVI Response
Inclusion of additional treatment effects: add triple therapy, sarilumab, and baricitinib to the model	Update both the systematic literature review (SLR) and network meta-analysis to ensure that the model reflects the most recent clinical evidence available.
Expand model functionality	Allow for an increase in model flexibility by (i) adding more utility functions, (ii) allowing for treatment specific discontinuation, (iii) including additional outcomes correlated with HAQ based on literature (or Corrona data) and (iv) adding biosimilars as a treatment option (with a potential discount rate)
Incorporate Corrona data (real-world, observational data) to update the model	Utilize Corrona data to model the impact of treatment on long-run HAQ progression, the size of HAQ rebound, treatment specific discontinuation, and potentially the effect of changes in HAQ and other outcomes of interest such as nursing home outcomes and fractures
More accurately capture treatment heterogeneity	YODA data from Yale University to measure whether treatment efficacy varies by number of prior tDMARDs, patient characteristics, or type of prior tDMARD received



Anticipated Timeline for All Revisions

Systematic Literature Review Update	July - September 2018
Investigate Availability of Yoda Data	August – September 2018
Submission to Corrona for Potential Use of RWD	September – October 2018
Network Meta-Analysis Update	December 2018 – February 2019
Update Model Functionality	January – March 2019
Update Value Tool and Expert Interface	April – May 2019
Release of IVI-RA Model 2.0	May – June 2019
RA-Model Version 2.0 Public Comment Period	June – July 2019



Appendix A. Public Comments on Initial Release of IVI-RA Modeling Platform

Contents of Appendix include formal comments provided by:

- AbbVie
- RheumPac, American College of Rheumatology
- Arthritis Foundation
- Bristol-Myers Squibb
- Genentech
- ICER
- Institute for Patient Access
- Janssen
- Pfizer
- Sanofi Genzyme Regeneron
- Tufts University



Darius Lakdawalla & Jason Shafrin
Executive Director
Innovation and Value Initiative
11100 Santa Monica Blvd, Ste 500
Los Angeles, CA 90025

January 17, 2018

RE: Innovation and Value Initiative (IVI) Rheumatoid Arthritis (RA) Model – Response to Request for Public Comments,


Dear Dr. Lakdawalla and Dr. Shafrin,

Thank you for the opportunity to provide input into the IVI RA Value Model. As a strategic advisor, we have reviewed the current version of the RA open source model and have consolidated key feedback that we believe will further enhance this model. The feedback is organized into two key sections: model structure and model results.

Model Structure

1. The ability to choose one of three possible model structures in the IVI-RA Model Interface allows the user a good deal of freedom, however, it is difficult to identify which of the model structures was used in the IVI-RA Value Tool:
 - a. In fact, the IVI-RA Value Tool severely limits the options which previously were available in the IVI-RA Model Interface.
 - b. It would be helpful to have an appendix available which lists all of the default options which were selected and what the input values used in the Value Tool were utilized
 - c. A footnote explaining the modeling approach used would be very helpful.
2. While the Value Tool allows the user to switch comparator sequences, it is unclear how the default sequence is chosen, and whether it is the most relevant sequence to compare.
 - a. A description of this default population would be helpful. Assuming this is high disease activity with average of 3.28 prior csDMARDs. Based on defaults in the tool, appears the biologic sequence is being used first line without prior csDMARDs.
3. When selecting base-case settings for sequencing, the language could be somewhat confusing since it first discusses a lifetime time horizon, then says “....when they stop working.”

Reader may think the model refers to the patient ending employment rather than the medicine stop working. Also, it is not accurate to imply that everyone's medicine will stop working. We have patients on the same treatment for many years.




4. The model appears to operate with two treatment courses i.e. sequence 1, and sequence 2. It is not intuitively clear what “sequence 1” really is since there’s just a descriptor of “cDMARDs”. It would be helpful to know if “cDMARDs” is a series of commonly prescribed cDMARDs that are used in some rational order, or if it is just a bucket of cDMARDs.
5. In the sequence 1 vs 2 description, it would be helpful to understand clearly if patients are only included in sequence #2 after they had failed Sequence #1, and if they did have to fail #1 first did they fail just one cDMARD or multiple cDMARDs?

Results

1. In the clinical economic outcomes from treatment, it would be helpful to have a graph corresponding to total QALYs
2. Unclear how the incremental costs are calculated in the 'Value' tab. For example, the mean total healthcare costs are \$56.31k for Sequence 1 and \$557.88k for Sequence 2. Based on that graph, one would think the incremental costs would be \$501.57k. However, in the 'Value' tab it calculates the incremental costs as \$373.61k, it is unclear what accounts for this difference. Might be helpful in the Cost Effectiveness table to add in the actual QALYs and Costs for each sequence.
3. There is a wide variety in the ICER results corresponding to the exact same model inputs. This could be due to a small number of iterations chosen to reduce run-time. Having such a wide variation in ICER results could result in "cherry-picking" where users run the model multiple times and then report the set of results they find most favorable.
4. The default Willingness to pay per QALY of \$150,000 seems arbitrary and very high, and there are no references to support this figure. The base case result would only be considered cost-effective using such high Willingness to pay per QALY, and thus the results may be misleading.
5. The MCDA piece is an interesting wrinkle. The weights themselves are reasonably intuitive, however, how the actual MCDA values are actually calculated are somewhat opaque.
 - a. Would be helpful to provide additional details on how the weights for this base case scenario were established. Is this based on patient-focused research?

Yours Sincerely,



Jenny Griffith, Pharm.D.
Associate Director
Health Economics and Outcomes Research
AbbVie

Jerry Clewett, Pharm.D., MBA
U.S. Scientific Director
Healthcare Solutions
AbbVie



Harry Gewanter, MD

Received via email

Hi there!

I like your model very much as it provides more depth and substance to the decision-making process by incorporating factors other than just money. I do have a number of suggestions/comments.

- I realize this is a near-impossible task, but the problem with this (and all other models) is that what you consider “cost” is really “list-price”. Nobody knows what the real costs are given the lack of transparency within the the health-care systems. We may know AWP, but we do not know the discounts, rebates, other fees and monetary shenanigans utilized by insurers, pharmacy benefit managers, pharmaceutical manufacturers or distributors that constitute the true costs. Further, very few physicians can tell you how much it “costs” to see an individual patient. I understand this is a systemic problem and therefore your model is comparable with, say, ICER as y’all are presumably using the same data, but I think you may want to put in a disclaimer about what is truly being evaluated.

- There are a number of areas where I think you can improve the explanations of your findings.
 - How much change in the HAQ is significant? What is the context for the numbers (ie, what does a “2” mean)?
 - Adding a ruler or better measurement benchmarks to the Life Expectancy (and other) graphs would make them more meaningful and understandable. In your standard example, I couldn’t suss out how much difference was present.
 - I think it would be helpful to include “denominators” and better/clearer descriptors for your various findings. For example, are the Healthcare Costs listed per year, per lifetime, per decade, etc? Are these for the individual or the population? The same can be said for the Days in Hospital, Earnings, Number of Infections, etc.

- While your glossary is helpful, it needs to be made into more readable English and less “Economic-Ese”. Having specific explanations and examples for non-economists would make this project more user-friendly, understandable and acceptable. Along the same line, since much of the model is based on QALYs and “Willingness to Pay” criteria, more robust explanations of these terms are necessary.

- Where did the \$150K/QALY come from? Knowing the source would make it more believable as a baseline figure.

- Within your MCDA page, the slider scales are not intuitive and I was not sure how to interpret what it meant when I moved the slider to the left or the right. Are these integral scales, logarithmic or something else? Again, more context is necessary to make this valuable aspect of the model more useful and practical.

- I think it would be helpful to try to identify additional patient preference issues and add them to the model. For example, does the age of the patient matter? What about an upcoming significant event, such as the birth of a grandchild? We know those issues change one’s “value system” regarding health care choice; I don’t know how to include them in the model, but I don’t see a place for these critical aspects of decision-making.



- I realize these are extraordinarily difficult components to include, but consideration of which joints are involved (ie, hands vs hips vs knees vs jaws, etc) and the impact of surgery on the disease, productivity, etc would make the model even more “realistic”. Similarly, there are a number of “extra-articular” issues that go along with RA beyond infections (ie, uveitis, rheumatoid lung, etc) as well as “extra-articular” benefits of treatment (ie, decreased cardiovascular disease) that could be added into the model as they are confounders.

- I loved having the productivity measure within the model. Treating RA is not just a “debit” of the treatment costs, it is also a “debit” of loss of income and community participation. However, along this thread, it might be helpful to add a similar “family” community slider (if that is possible) since someone with RA may result in additional family/community care taking costs, both direct and indirect and these are not small amounts.

- I liked the concept of what it meant to have RA if you are healthy. Not something usually considered.

- I don’t know how hard it would be to add more options to the “Not entirely average” modifier (ie, quintiles, etc) to allow additional explorations of response.

I hope these thoughts are helpful. Keep up the good work and keep improving the model.

Harry L. Gewanter, MD, FAAP, MACR

Medical Director, Medical Home Plus, Inc
Chair, RheumPAC, American College of Rheumatology
Associate Clinical Professor of Pediatrics, Children’s Hospital of Richmond at VCU

hgewanter@icloud.com
(C) 804.307.6896

March 30, 2018

Innovation and Value Initiative
Open Source Value Project: RA Model
Public Comment

Jennifer Bright, MPA
Executive Director

Mark Linthicum, MPP
Director of Scientific Communications

The Arthritis Foundation appreciates the opportunity to provide comments on IVI's Open Source Value Project RA Model. Below please find our team feedback as well as the collective feedback from the review conducted at our Patient Leadership Council meeting on March 11, 2018.

General Feedback

The tool is somewhat difficult to use/understand and, depending on the specific function/parameter can be either too complex or too simple. That said, it seems like it could have potential, particularly if it was nimble enough to be used in real time (i.e., imagine a Congressional briefing where a PAO can show the longitudinal cost effectiveness of a public health program paying for a specific category of drugs). One key challenge is that the tool is incredibly complicated (in its current form) for a patient or PAO to use, or perhaps anyone who isn't an HEOR expert, but it seems too simple for an actual HEOR expert to be using (or someone on the "inside" of a health plan/insurer who would be relying on their own actuaries). Might it be necessary to further customize the tool depending on the intended end-user?

Specific comments:

- The purpose of the tool as stated is confusing/concerning; on the opening frame there is no mention of identifying treatments (even at a population level) that are likely to have the best treatment effect. Rather, the only stated goal relates to allocating healthcare dollars. It seems like the portion of the equation dedicated to working to improve health outcomes for patients should be given some above-the-fold space?
- It is great that two patient focus groups were conducted but that doesn't seem to be a robust enough patient engagement approach. What was the role of the patients on the steering committee and can you share any examples of the types of insights patients shared that actually shaped the tool? A more robust, iterative patient engagement approach would likely be fruitful— one that begins at the very earliest stages of conceptualizing a disease-specific value tool.
- The treatment sequence information is difficult to follow; is the suggestion that all of these drugs were used? For how long? What was the impetus for d/c and beginning a new one? Was there a lag time between treatments due to access challenges (step therapy, prior auth., etc.)?

- The time parameter/amount of fluctuation that occurs but is not captured or visualized in the “Change of patient functional status” bars is confusing. It is nice to see in vivid color the significant difference between HAQ scores in DMARD-only treatment vs. biologic treatments, but with such a volatile disease, seeing a static representation of what happens over a lifetime seems incomplete.
- Do the underlying assumptions in the model fully represent the patient population?
- Is there really enough data to know impact of biologics on life expectancy? People have only had access to them since 2000; are you including people who were diagnosed prior to that and treated with non-biological DMARD cocktails first, and then switched to biologics? If so, that may need to be teased out/explained.
- Given that this is an RA tool, why wasn't RAPID-3 used?
- What is the reasoning for the “value to the currently healthy” element of the model? It seems like this would be more relevant in contagious diseases?
- QALYs have pretty significant limitations from a disease/disability perspective and don't offer a full, humanized perspective of the real experience of RA. In the absence of another measure to use, it makes sense that QALYs were used, but as new options emerge, they should be taken into consideration.

Arthritis Foundation Patient Leadership Council Review (3.11.18)

General Feedback

- Does not seem very user-friendly for patients.
- Calculating based on these particular data is concerning because it seems to fly in the face of current thinking around genetic markers and other medical predictors of treatment benefit.
- What might be valuable for an individual user is the ability to enter my own sequence, and some other key demographic factors (age, sex, length of diagnosis) and see what the comparative value is of the next medication options.
- If it is truly for patients, it needs to be more usable and accessible to patients; if it is really intended for population level analytics and HEOR, then that's a very different animal.

Model-Specific Feedback

Opening Screen/Welcome to the IVI RA Tool/Get Started

Comments:

- “How to allocate limited healthcare dollars” is a worrisome sentence. It seems that the focus is only on the spending side and not on the treatment and health outcomes side.
- Who is funding this tool? If pharmaceutical companies are funding it, it seems like it would be in their best interest to promote access to drugs but the tone here seems to be limiting access.
- We are already fighting for access to drugs on so many levels. I understand that there are “limited healthcare dollars” and maybe it is just the way they've phrased it but it automatically indicates someone isn't going to get what they need (who's going to “get it” and who's not going to “get it” and who is determining that?)

- I don't know that I want population-level analysis defining what MY treatment is. I'm afraid that the tool will result in insurance companies using population-level standards as the criteria for treatment.
- It's vague; it isn't clear what is meant by "limited healthcare dollars"—is that referring to practices of insurance companies? Something else?

RA Treatment History

Comments:

- Some patients will not know what a DMARD or biologic is
- Is Xeljanz considered a biologic in this tool?

Gender Ratio

Comments:

- A great deal of confusion on this piece; why would we be able to toggle gender? Isn't this static? Why should people be able to change it?
- Confusion about the apparent discord between whether this can be used for individuals or is it strictly for population-level analysis.
- Again—who is this tool for? Is it to be used by individuals, physicians, insurers, etc.? (Why would an individual patient care about the gender ratio of a population—and how would they know?)

Age

Comments:

- Is this so that when an individual is using the tool they can toggle to their age?
- Same questions/issues here as with gender—unclear why we should be able to move this.

Sequence of RA Treatments

Comments:

- Treatment sequence is so incredibly variable and dictated by several factors like insurance; what assumptions were made/what data was used to generate the default sequences?
- If this is for patients to use, should use brand names for easier recognition
- What if you don't take methotrexate with your biologic? Many patients do not take MTX—and either take another DMARD (Arava, Azathioprine, etc.) or no concomitant DMARD.
- Why is Humira the first in the default sequence (and why are the others listed in the order that they are?) Is there data to support that it is often the first sequence? What are the data sources? (Were patients asked?)
- What about breaks in treatment for pregnancy, illness, step-therapy? There is no way to show pauses/breaks in the sequences.
- What about returning to previously tried drugs—which also happens? Is there a way to build repeats into the sequence?
- Where is Remicade?
- Your insurance often dictates what you start with; on Medicare it might be Remicade versus ESI which might require Enbrel or Humira first. How are the sequences reflecting the fact that the "choice" isn't necessarily a "choice"?

Treatment Cost

Comments:

- Costs to whom? Are these OOP?

Change in functional status/HAQ score

Comments:

- Result resonated—standard DMARD therapy less effective than biologic sequence

Life expectancy

Comments:

- Results are plausible, but many questions about who is in which population
- Does not seem like we are really comparing apples to apples
- Generally speaking, no one is allowed to “start” on a biologic—you must have been on a DMARD first (under current insurance restrictions) so how long did people in category 2 wait before they started a biologic? How much damage was done before a person began biologic therapy?

Total Healthcare Costs

Comments:

- Mixed reactions because cost experience has varied
- Concern that a more newly diagnosed person accessing this tool might be scared away from biologics
- Costs “to whom”? Are these OOP?
- What it tells me if I’m a health insurer, is that I want to keep people on DMARDs as long as I can because length of life is roughly the same but DMARDs are cheaper (which is a concern for patients)
- Sequence 1 costs seem too low—it does not look like productivity, disability, etc. are factored in?
- Same issue here with who is in which pot—who is in the biologic group? How long did they wait to get on a biologic? Are there people who were diagnosed many years ago and had to wait 10-15 years for biologics?

Number of Days in Hospital

Comments:

- It isn’t clear what the causes are here. Are these due to side effects from treatment or exacerbations of disease because of ineffective treatment? Joint replacement complications?
- Concerns about the standard deviation here—aren’t results inconclusive?

Reductions in Patient Earnings

Comments:

- Interesting that this is captured here and not under “total costs”

Number of Serious Infections

Comments:

- How did the data parse out complication of reaction to infection being masked by efficacy of the biologic?

- Applies to apples concern here; biologics range by mechanism and how targeted they are so lumping them into one bucket seems unhelpful
- Another standard deviation issue here—if the SD is the entire population than the sample must not have been appropriate

Cost-effectiveness Table/Graphs

Comments:

- Confused why everything in sequence 1 is “non-applicable” in our particular run
- Confusion around the use of “lifetime” in the model; if biologics have only been around for 20 years (and some far less) what constitutes a life time?

MCDA Weights

Comment:

- Why is default “no value” given to mode of administration (and additional challenge here—can’t assume infusion is “least desirable”—some people prefer because it is what insurance dictates and/or because they prefer to not self-administer)
- Why is default “no value” given to years since FDA approval?
- A few things missing:
- Fatigue
- Side effects—not “serious” events, but those frequent bothersome issues that impact QOL

We look forward to seeing future iterations of the tool.

Sincerely,

M. Suzanne Schrandt, JD



Director, Patient Engagement
Arthritis Foundation



U.S. Pharmaceuticals
P. O. Box 5326 Princeton, NJ 08543-5326

January 17, 2018

Darius Lakdawalla & Jason Shafrin
Innovation & Value Initiative
11100 Santa Monica Blvd
Los Angeles, CA 90025
research@thevalueinitiative.org

BY ELECTRONIC DELIVERY

Re: Proposed IVI-RA Model

Dear Dr. Lakdawalla and Dr. Shafrin:

As a leader in immunology research, Bristol-Myers Squibb (BMS) acknowledges the importance of understanding and fully characterizing the value that innovative therapies provide to patients, and appreciates the opportunity to comment on the Innovation and Value Initiative (IVI) Rheumatoid Arthritis (RA) Model as part of IVI's Open Source Value Project (OSVP). BMS is dedicated to advancing the science of immunology and to disseminating the results of our research to ensure that our work can benefit the widest range of patients.

We have reviewed the current version of the IVI-RA model and believe it could benefit from the following feedback to guide its ongoing development.

Refine with HLC's Principles of Value Frameworks

BMS agrees with IVI's patient centric and transparent approach to developing the current version of the IVI-RA model. For example, the patient focus groups conducted initially were a key resource in developing the model. As IVI looks to refine the RA model further, we recommend referring to a set of principles on value frameworks developed by the Healthcare Leadership Council (HLC), a coalition of chief executives from all disciplines within American healthcare. In 2017, HLC released the following set of principles that should guide the creation of value frameworks being used to determine the cost-effectiveness of new healthcare innovations¹. These align also with our own company's principles on value frameworks.

1. Collect patient and provider input on what "value" should be measured in a treatment option in order to measure outcomes that matter to patients and providers (such subjective data as discomfort during or after treatment).
2. A diverse group of disease area experts should participate in both the development of methodology and assessments before these are submitted through a peer review process to ensure scientific rigor.
3. During and after the review process, provide full transparency of evaluation criteria, including any models and data used – allowing for research to be analyzed and results replicated by others.

4. Assess and reassess value over time (recognizing that prices can vary over time), to capture appropriately the value of curative or preventative treatments whose full value may not be realized until after the initial approval.
5. Define value to society broadly – this includes outcomes values such as productivity, opportunity costs, and avoided long-term costs.
6. Incorporate real-world evidence and adjust evaluation techniques to capture actual patient outcomes and preference for treatment, recognizing some data may be difficult to obtain for pragmatic or ethical reasons.
7. Treatments should accurately reflect real-world usage.
8. Consider variations in treatment setting, technique, and provider when evaluating a new product or technique.

Already with the beta version of the RA model, IVI has adhered to many of these principles. As IVI seeks to refine the model, we recommend robust patient and provider engagement, as well as increasing the incorporation of real world evidence. With respect to capturing patient value, we recommend referring to perspectives from the advocacy community in a recent publication in Health Expectations for guidance on additional IVI engagement with patients and patient groups.ⁱⁱ

Expand the Model to More Fully Capture All Elements of Value Rather than a Focus on Quality-Adjusted Life-Years (QALYs)

We are encouraged IVI has incorporated other value elements other than the QALY; however, the beta version of IVI-RA proposes the primary health outcome is the QALY. Experts have identified other value elements and BMS believes rather than promulgating the use of this limited measure, value frameworks should aim to incorporate as many elements of value as possible. For example, ISPOR's Special Task Force (STF) on Value Assessment Frameworks has identified twelve different components of value in their draft whitepaper – including QALY, net costs, productivity, adherence improving factors, reduction in uncertainty, fear of contagion, insurance value, severity of disease, value of hope, real option value, equity, and scientific spillovers. This catalogue of elements provides one of the first, truly forward-looking steps in comprehensively characterizing all of the multidimensional facets of value. We recommend prioritizing the incorporation of additional value elements into the IVI-RA model in the revision process. While we appreciate that not all of these value elements can be incorporated in every model, as appropriate to the disease state they could further help estimate value. Not only will this help advance the methodology for the value elements identified by ISPOR's STF, but it will more fully capture value to patients and society.

Expand the Role of Safety in the Model and Differentiate Safety Outcomes

In addition to inadequate efficacy, lack of safety in the form of adverse events is also a common reason for RA patients switching medications. The model's primary focus is on patients who switch or discontinue their medication due to efficacy only. This presents a bias toward products with a better efficacy profile and worse safety profile. The model allows switch due to adverse events, but only in the case of serious infection, as these “have a significant cost impact and increased risk over background rates to be meaningful to include.” Additional

adverse events are important to patients and result in treatment switching. Therefore, the model should expand beyond serious infections and include other adverse events.

The model also assumes that the infection rate is equal among targeted disease-modifying antirheumatic drugs (tDMARDs) because “the published results for specific tDMARDs are estimated with very little precision.” Evidence shows that specific tDMARDs have decreased infection risk compared to other tDMARDs. In RA patients with prior exposure to a biologic agent, exposure to etanercept, infliximab, or rituximab was associated with a greater 1-year risk of hospitalized infection compared to the risk associated with exposure to abatacept.ⁱⁱⁱ Among RA patients who experienced a hospitalized infection while on anti-TNF therapy, abatacept and etanercept were associated with the lowest risk of subsequent infection compared to other biologic therapies.^{iv} The expected costs of serious infections were lower for IV and SC abatacept than for infliximab and adalimumab in hypothetical analyses based on two large clinical trials.^v Therefore, the assumption of equal infection rates negatively impacts the performance of these tDMARDs in the model. The model should differentiate rates of serious infection among tDMARDs.

Allow for Analysis Based on Poor Prognostic Factors

Although this model allows more patient heterogeneity than previous cost-effectiveness models, there is still no way to analyze subgroups, such as those with poor prognostic factors and more severe disease. Even if this data is not available across every clinical trial in the network meta-analysis (NMA), the final tool should allow the user to conduct a scenario analysis for different subgroups.

Select the Most Appropriate Outcomes

The model relies on each agent’s impact on Health Assessment Questionnaire (HAQ) Disability Index score during the initial treatment phase and incorporates parameters such as progression defined by HAQ over time, mortality rates as a function of HAQ-based progression, quality of life based on HAQ, and cost based on HAQ-progression. Focusing on this efficacy measure during the initial treatment phase, and no other outcomes, is a limitation. The model should seek out ways to incorporate other measures such as Disease Activity Score (DAS) and American College of Rheumatology (ACR) response (ACR20, ACR50, ACR70, ACR90) to define model outcomes, even if evidence is not as robust as that for HAQ.

Adjustments for HAQ progression for sequencing of patients

BMS would like to bring to IVI’s attention that the reference Wolfe and Michaud (2010) for the model’s HAQ progression does not take into account the patients’ treatment history.^{vi} Specifically, a subgroup of patients have been shown to benefit from a TNF to non-TNF sequence compared to TNF cycling.^{vii}

Bristol-Myers Squibb appreciates IVI’s efforts to engage stakeholders in the development of the IVI-RA model and we look forward to providing continued input to IVI as it refines the model. We welcome the opportunity to meet to further discuss our feedback. If you have any questions, please do not hesitate to contact Michael Ryan, PharmD, Head of U.S. Value, Access &



U.S. Pharmaceuticals
P. O. Box 5326 Princeton, NJ 08543-5326

Payment at (609)302-3198 and Mitch Higashi, PhD, Head of U.S. Health Economics Outcomes Research at (609)302-3798.

Sincerely,

Michael Ryan, PharmD
Head of U.S. Value, Access & Payment

Mitch Higashi, PhD
Head of U.S. Health Economics & Outcomes Research

References:

-
- ⁱ Healthcare Leadership Council. Principles on Value Assessment Frameworks. Available at: <https://www.hlc.org/app/uploads/2017/05/HLC-Final-Principles-on-Value-Frameworks.pdf>
- ⁱⁱ Addario BJ, Fadich A, Fox J et al. Patient value: perspectives from the advocacy community. *Health Expectations*. 2017;1-7. Bristol-Myers Squibb provided funding to support the writing and editing of this paper and provided a grant to ensure the paper was published with Open Access.
- ⁱⁱⁱ Yun H, Xie F, Delzell E, Levitan EB, Chen L, Lewis JD, Saag KG, Beukelman T, Winthrop KL, Baddley JW, Curtis JR. Comparative Risk of Hospitalized Infection Associated With Biologic Agents in Rheumatoid Arthritis Patients Enrolled in Medicare. *Arthritis Rheumatol* 2016, 68(1):56-66.
- ^{iv} Yun H, Xie F, Delzell E, Chen L, Levitan EB, Lewis JD, Saag KG, Beukelman T, Winthrop K, Baddley JW, Curtis JR. Risk of hospitalized infection in rheumatoid arthritis patients receiving biologics following a previous infection while on treatment with anti-TNF therapy. *Ann Rheum Dis* 2015, 74(6):1065-71.
- ^v S Johnston, S Kelly, A Nadkarni, KL Wilson, B Limone, M Hochberg. Healthcare Costs Associated with Serious Infections Among Biologic-Naïve Rheumatoid Arthritis Patients Initiating First-Line Biologic Treatment. ACR/ARHP Annual Scientific Meeting, November 14-19, 2014, Boston, MA.
- ^{vi} Wolfe F, Michaud K. The loss of health status in rheumatoid arthritis and the effect of biologic therapy: a longitudinal observational study. *Arthritis Research & Therapy* 2010, 12:R35.
- ^{vii} Gottenberg J-E, Olivier B, Perdriger A et al. Non-TNF-Targeted Biologic vs a Second Anti-TNF Drug to Treat Rheumatoid Arthritis in Patients with Insufficient Response to a First Anti-TNF Drug: A Randomized Clinical Trial. *JAMA* 2016, 316(11):1172-1180.



Bristol-Myers Squibb

Received via email from Alexander Marshall

Hi Jason and Devin,

Thank you for taking the time to speak with us today on the model. Here is the citation for the AMPLE article we discussed. We also wanted to follow up on key points from the meeting and next steps.

Schiff M, Weinblatt ME, Valente R, et al. Head-to-head comparison of subcutaneous abatacept versus adalimumab for rheumatoid arthritis: two-year efficacy and safety findings from AMPLE trial. *Ann Rheum Dis*. 2014 Jan;73(1):86-94.

Since the package insert states the abatacept IV loading dose is optional and the pivotal head to head RCT AMPLE does not include an IV loading dose:

- We agree with the suggestion to have the default option for abatacept SC without an IV loading dose.
- When the option to include a loading dose is chosen by the user, it should include a reduced infusion cost. Currently, the infusion cost is the same between abatacept IV and abatacept SC. It should be lower for the SC formulation as there is only one infusion associated with this regimen.

For next steps regarding medications that are available in both IV and SC formulations (Orencia, Actemra, and Simponi):

- Please let us know whether the SC or IV form of golimumab and tocilizumab are currently being used in the model.
- We believe the model should be transparent as to the formulations used for all drugs. Therefore, we recommend all drugs are labeled by their formulation and that all formulations currently available for RA are included in the model.

Thank you again and we look forward to hearing from you!

Best Regards,

Alex and the BMS Team



February 16, 2018

Innovation and Value Initiative
11100 Santa Monica Blvd, Ste 500
Los Angeles, CA 90025

RE: Public Comment on the Open Source Development of the IVI-RA Model

Dear IVI-RA Technical Expert Panel:

Genentech, a member of the Roche Group, appreciates the opportunity to provide comments on the Innovation and Value Initiative's (IVI) rheumatoid arthritis (RA) model. We commend IVI's efforts in facilitating an open source, transparent process for estimating the value of health care technologies for a broad set of stakeholders.

Genentech is committed to advancing the scientific understanding of RA and pursuing the development of novel therapies to help individuals diagnosed with this chronic and debilitating disease. Treatment decisions for patients with RA are complex and personal. We support initiatives that account for the needs of individual patients and facilitate a meaningful dialogue between a broad set of stakeholders.

Our comments are focused on enhancing IVI-RA model's usability for a diverse set of users and clarifying model assumptions. We hope our recommendations contribute to improving the IVI-RA model and advancing the field of value assessment.

IVI-RA Value Tool

Additional enhancements to the model will increase the end user's ability to customize it to meet their individual needs and obtain information most relevant to their decision making. For those unfamiliar with the complexity of the RA disease landscape, the numerous customization options risk confusion. Additional context and guidance will better equip users in navigating the model and obtain meaningful analyses. Genentech recommends the following:

1. Avoid potentially biasing the user by removing the default sequence in the Setup function of the model. The model default currently compares a sequence of six "targeted" DMARDs to conventional DMARDs alone. Multiple, alternative treatment sequences are

consistent with ACR guidelines and real-world practice due to the heterogeneity and complexity of RA. The ACR guidelines recommend that patients with established RA who fail DMARD monotherapy move to combination DMARD therapy, add a necrosis factor inhibitor (TNFi) with or without methotrexate (MTX) or add a non-TNFi biologic with or without MTX, in no particular order of preference.¹ The guidelines also conditionally recommend using a non-TNFi or a TNFi after TNFi failure. The default sequence currently in the model begins with two TNFis in combination with MTX, i.e., adalimumab followed by etanercept. However, clinical evidence supports switching to a non-TNFi after first TNFi failure.^{2,3} Therefore, since many alternative options consistent with ACR guidelines are available, we recommend removing the default sequence, and allow users to choose their own sequence.

2. Mitigate potential misinterpretation of health care costs presented in the Outcomes function by clarifying that the costs represent those incurred over the lifetime model time horizon, which include costs after completing treatment(s) specified in model comparators.
3. Enable greater customization to user perspective by including the ability to modify the model time horizon since preferred time horizons will vary based on user perspective.
4. Facilitate ease of use of the model by describing source data that informs the clinical and economic outcomes for each treatment option in the Outcomes function.
5. Guide users to a more accurate interpretation of the Health Assessment Questionnaire (HAQ) Disability Index results, an important measure of patients' health status in RA, by including alternative means of presenting the change in functional status, such as "better" and "worse" on the y-axis.

IVI-RA Model Interface

The IVI-RA Model Interface is provided as an option for users who want increased control over model parameters and structural assumptions. We appreciate IVI for developing a highly flexible model. To increase ease of use for those seeking greater customization of the IVI-RA model, we recommend the following:

1. Guide users with pertinent details upfront in the introduction in order to avoid confusion. For example, we recommend providing a diagram to detail the IVI-RA model is a discrete-time individual patient simulation that simulates outcomes for individual patients.
2. Provide additional background and implications around the selection of the HAQ, ACR and DAS28 in respect to treatment and treatment switch.

3. Enable users to have a sound understanding of their starting point by outlining the rationale for base case assumptions in the Model Interface. For example, it is not readily apparent why a generalized gamma distribution for treatment duration was chosen.
4. Comprehensively account for uncertainty by including a deterministic sensitivity analysis and complementing the existing probabilistic sensitivity analysis.

Multi-Criteria Decision Analysis (MCDA)

We commend IVI for incorporating MCDA into IVI-RA given traditional measures of value, such as cost-effectiveness analyses, may not adequately capture value for all health care stakeholders. It is important that treatment attributes most valuable to patients are integrated into health care decision making. As stated by IVI, we agree that factors such as patient out-of-pocket costs and side effect profiles should be included in the MCDA framework and highlights an important next step for IVI-RA. In addition, we recommend providing background information to how best to interpret the MCDA with direct links to the technical report and additional information boxes. Furthermore, clarification around how MCDA parameters were placed on a common scale and the rationale for default settings will help users navigate and customize the MCDA to meet their needs.

Additional Recommendations

We recommend the following corrections and modifications to accurately reflect current available treatment options in RA.

1. Tocilizumab monotherapy, in subcutaneous (SC) and intravenous (IV) formulations, should be included as treatment options based on its FDA approved indication⁴. It is important to include both SC and IV formulations given the importance of treatment administration in the MCDA.
2. Treatment costs (Set-Up function) should account for different dosing and administration options available for tocilizumab, which includes SC weekly and IV⁴. Currently the IVI model specifies every other week only.
3. Biologic DMARDs should be consistently referred as “bDMARDs” in order to avoid potential confusion with tDMARDs. tDMARDs is sometimes used to denote traditional DMARDs.
4. Rituximab dosing should be consistent with prescribing information. Rituximab IV dosing after two initial infusions is every 24 weeks or based on clinical evaluation, but not sooner than every 16 weeks⁵. Note that rituximab monotherapy is considered off-label as it is FDA-approved to be used in combination with MTX.

Genentech offers IVI our support and partnership given our long standing position in RA and desire to advance the field of value assessment. We hope that IVI finds our comments helpful in the next iteration of the IVI-RA and achieving a transparent process for estimating the value of health care technologies for diverse health care stakeholders.

Sincerely,

A handwritten signature in blue ink that reads "Jan Hansen". The signature is written in a cursive, flowing style.

Jan Hansen, PhD
Vice President, Evidence for Access
Genentech U.S. Medical Affairs

References:

1. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)*. 2016 Jan;68(1):125. <https://www.rheumatology.org/Portals/0/Files/ACR%202015%20RA%20Guideline.pdf>
2. Gottenberg JE, Brocq O, Perdriger A, et al. Non-TNF-targeted biologic vs a second anti-TNF drug to treat rheumatoid arthritis in patients with insufficient response to a first anti-TNF drug: A randomized clinical trial. *JAMA*. 2016 Sep 20;316(11):1172-1180. <https://www.ncbi.nlm.nih.gov/pubmed/27654603>
3. Emery P, Keystone E, Tony HP, et al. IL-6 receptor inhibition with tocilizumab improves treatment outcomes in patients with rheumatoid arthritis refractory to anti-tumour necrosis factor biologicals: results from a 24-week multicentre randomised placebo-controlled trial. *Ann Rheum Dis*. 2008 Nov;67(11):1516-23. <https://www.ncbi.nlm.nih.gov/pubmed/18625622>
4. Actemra (tocilizumab) [Package Insert]. Genentech Inc.; South San Francisco, CA; 2017. https://www.gene.com/download/pdf/actemra_prescribing.pdf
5. Rituxan (rituximab) [Package Insert]. Genentech Inc.; South San Francisco, CA; 2015. https://www.gene.com/download/pdf/rituxan_prescribing.pdf

January 17th, 2018

Innovation and Value Initiative

11100 Santa Monica Blvd, Ste 500

Los Angeles, CA 90025

Dear IVI-RA modeling team,

Thank you for the opportunity to provide feedback on the RA-IVI model. A team at the Institute for Clinical and Economic Review (ICER) has reviewed the model methodologies, R source code and online model interface. Please find below our comments, suggested revisions, and associated references for your consideration.

Sincerely,

Varun Kumar

Health Economist

Institute for Clinical and Economic Review

Two Liberty Square, 9th Floor

Boston, MA 02109

617-528-4013 x7011

vkumar@icer-review.org

www.icer-review.org

Major

1. Formulate treatment switching rules (S1-S6) as part of the decision rather than an uncertainty. In section 2.4, one stated use of the model is for evaluating consequences of clinical guidelines. For example, users can (in theory) examine different treatment switching decision rules. However, in the current model setup, treatment switching is considered a "structural uncertainty" rather than a decision or strategy in S1-S6 (although it is symbolized as a decision in Figure 3). It is also unclear why the choice for modeling changes in HAQ scores (H1-H3) limits the available switching strategies. For example, why does modeling HAQ directly preclude treatment switching based on ACR? This limitation has implications throughout the document and the model.
2. Elaborate on the reasons for the differences in results when using the different approaches for modeling HAQ in the initial treatment phase (H1-H3). Were any parameters calibrated to match known scenarios? The data for ACR to EULAR were based on US veterans, which may not be representative of the overall RA population in the US. The other studies similarly may not be representative of the target population.
3. Elaborate on justification for using observational data by Wolfe and Michaud 2010 for deriving annual HAQ change in the 'constant linear rate of progression' approach. ICER recommends using the National Data Bank (NDB) for Rheumatic Diseases dataset in NICE DSU 2015 (Progression of Disease in People with Rheumatoid Arthritis Treated with Non Biologic Therapies), which more closely resembles the model's target population based on gender and age, although HAQ scores in both datasets are similar.¹ The mean annual HAQ change reported in the NICE DSU for the NDB population is 0.0269 while the RA IVI model uses a 0.031 annual HAQ change. While the Michaud et al. 2011 study that IVI has used provides HAQ by age group as well as HAQ based on duration of disease, it does not state duration of disease by age group. The linear relationship of HAQ change over time is not known beyond length of registry follow-up data, about 15 years on average, and may likely flatten after. ICER recommends using a constant HAQ degradation beyond 15 years in the model. Thus, associating a HAQ change by age group without equating to duration of disease may misrepresent actual HAQ change over time. We recommend including a HAQ trajectory over the lifetime of the model in graphical or tabular format for cDMARDs and default tDMARD sequence.
4. ICER recommends including the independent effects of radiographic progression, using (modified) total Sharp score (TSS) on HAQ. While its effect may be modest, it adds clinical face validity to the model.²⁻⁵
5. In alignment with including the TSS relationship with HAQ, ICER recommends using OR of 1.97 (in patients with radiographic data) as reported by Wolfe et al. 2003 for the effect of HAQ on mortality.⁶ This will result in log odds of 0.679. Please elaborate on calculations used to derive log HRs for a 0.25 unit- increase in HAQ, since the Michaud et al. 2012 study reports this unit increase annually after the first six months. Please provide justification on assuming the same log HR for a 0.25 unit-increase in HAQ after 36 months. An alternative approach to derive RA-specific mortality is to use the equation:

Mortality rate = Mortality from life table*1.33^{HAQ}, which has been used in another RA model.⁷

6. Elaborate on the rationale for assuming a normal distribution of HAQ rebound between 0.7 and 1 in the default settings. In the absence of evidence on HAQ rebound following treatment discontinuation, ICER recommends a conservative approach assuming 'rebound' to baseline HAQ when prior to subsequent treatment initiation.
7. Elaborate on the model validation process since section 10 is lacking details. Recommend providing details on all validation processes and results, following the ISPOR/SMDM recommendations.⁸

Minor

8. Revise notation throughout the document, including in the appendix, for consistency. For example, the use of "T" as a variable or to symbolize the transpose of a matrix, the use of commas in subscripts or not, the use of each Greek (and Latin) letter to represent a specific parameter rather than being used to represent different parameters in different equations, among others. The consistent use of notation would add clarity to the document.
9. Include an impact inventory for societal perspective, including the elements identified as important but not included due to lack of data or other limitations.
10. Revise Algorithm 1 for clarity and completeness. For example, there are several "if" statements without corresponding "else" statements.
11. In the online version, provide guidance on the appropriate distribution to use for time to treatment discontinuation. While allowing for 7 distributions enhances flexibility, it also adds confusion for users who cannot easily see how well the model fits the data.
12. The online simulation model is slow and prone to time out. If this version is to be implemented more widely in practice, we suggest improving the efficiency of the R coding, switching to a server faster than the current Rshiny server, or considering the use of meta-modeling/emulator techniques.
13. We suggest adding relevant citations in Section 1 to support current challenges with using and disseminating models.
14. We suggest including costs and probability of TB. While these estimates aren't key drivers of model results, they offer clinical face validity.
15. Use data available for most agents on specific rates of serious infections, rather than a summary rate.⁹⁻¹¹

References

1. Gibson L, Alava MH, A W. *Progression Of Disease In People With Rheumatoid Arthritis Treated With Non Biologic Therapies; Report By The Decision Support Unit*. Sheffield, UK: School of Health and Related Research, University of Sheffield;2015.
2. Stephens S, Botteman MF, Cifaldi MA, van Hout BA. Modelling the cost-effectiveness of combination therapy for early, rapidly progressing rheumatoid arthritis by simulating the reversible and irreversible effects of the disease. *BMJ Open*. 2015;5(6).
3. Barnabe C, Hazlewood G, Barr S, Martin L. Comparison of radiographic scoring methods in a cohort of RA patients treated with anti-TNF therapy. *Rheumatology (Oxford, England)*. 2012;51(5):878-881.
4. Breedveld FC, Weisman MH, Kavanaugh AF, et al. The PREMIER study: A multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis and rheumatism*. 2006;54(1):26-37.
5. Lillegraven S, Prince FH, Shadick NA, et al. Remission and radiographic outcome in rheumatoid arthritis: application of the 2011 ACR/EULAR remission criteria in an observational cohort. *Annals of the rheumatic diseases*. 2012;71(5):681-686.
6. Wolfe F, Michaud K, Gefeller O, Choi HK. Predicting mortality in patients with rheumatoid arthritis. *Arthritis and rheumatism*. 2003;48(6):1530-1542.
7. Carlson JJ, Ogale S, Dejonckheere F, Sullivan SD. Economic evaluation of tocilizumab monotherapy compared to adalimumab monotherapy in the treatment of severe active rheumatoid arthritis. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2015;18(2):173-179.
8. Eddy DM, Hollingworth W, Caro JJ, Tsevat J, McDonald KM, Wong JB. Model transparency and validation: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force-7. *Medical decision making : an international journal of the Society for Medical Decision Making*. 2012;32(5):733-743.
9. Smolen J, Genovese M, Takeuchi T. THU0166 Safety Profile of Baricitinib in Patients with Active RA: An Integrated Analysis. *Annals of the rheumatic diseases*. 2016;75:243-244.
10. Genovese MC, Fleischmann R, Kivitz AJ, et al. Sarilumab Plus Methotrexate in Patients With Active Rheumatoid Arthritis and Inadequate Response to Methotrexate: Results of a Phase III Study. *Arthritis & rheumatology (Hoboken, NJ)*. 2015;67(6):1424-1437.
11. Strand V, Ahadieh S, French J, et al. Systematic review and meta-analysis of serious infections with tofacitinib and biologic disease-modifying antirheumatic drug treatment in rheumatoid arthritis clinical trials. *Arthritis Research & Therapy*. 2015;17:362.



February 16, 2018

Submitted electronically to: research@thevalueinitiative.org

Darius Lakdawalla, PhD
Executive Director
11100 Santa Monica Blvd, Ste 500
Los Angeles, CA 90025

Re: Feedback on the Innovation and Value Initiative's Open-Source Value Project Model for Rheumatoid Arthritis

Dear Dr. Lakdawalla:

On behalf of the Institute for Patient Access, I thank you for the opportunity to provide feedback on the Innovation and Value Initiative's Open-Source Value Project model for rheumatoid arthritis.

About the Institute for Patient Access

The Institute for Patient Access (IfPA) is a physician-led policy research organization dedicated to maintaining the primacy of the physician-patient relationship in the provision of quality healthcare. To further that mission, IfPA produces educational materials and programming designed to promote informed discussion about patient access to approved therapies and appropriate clinical care.

IfPA was established in 2012 by the leadership of the Alliance for Patient Access, a national network of more than 800 physician advocates committed to patient access. IfPA is a 501(c)(3) public charity non-profit organization.

Approach to Value and Effectiveness Models

Too often, cost-effectiveness analyses exclude patients and their health care providers from the very calculations that determine whether they can access necessary diagnostics and drugs. Models that are shrouded in secrecy and findings that are conveyed in cumbersome, inaccessible language leave patients feeling overwhelmed and excluded.

IfPA is pleased to see that IVI has taken a different approach.

The open-source framework not only welcomes input from all stakeholders, but also allows them to customize measurements to reflect their own unique values. This far exceeds the usefulness of a static, inaccessible framework.

Comments on the Rheumatoid Arthritis Model

IfPA is pleased to see that IVI's rheumatoid arthritis model accommodates factors that matter to patients, such as whether a therapy is delivered orally or via IV, and the impact of different therapies on productivity. These are major factors in patients' experiences with a given therapy.

The model's level of customization is valuable for physicians and patients, who will be able to assess with a new level of precision how different therapeutic options will impact patients, and what costs and secondary factors make for the right choice. The model also acknowledges an important reality for rheumatoid patients and providers. A single therapy seldom proves effective for a patient's lifetime. Comparing a series of alternating therapies makes for a more realistic, reliable model.

As IVI works toward perfecting its model, developers may want to simplify the interface to account for users' varying levels of understanding about economic modeling. For instance, developers might consider visualizing results with icons and images that relate to the factor being quantified rather than using bar graphs.

It might also prove helpful to divide some steps into smaller sub-steps, allowing more context for each. For example, the page dedicated to "Clinical and Economic Outcomes for Patients" might be more accessible to patients if it were broken into functional status, life expectancy, and other components as individual pages. A summary page at the end of this section could synopsise the findings together.

Conclusions

IfPA commends IVI for working toward an open, customizable value framework that can be both useful and usable to a range of stakeholders, including the patients who feel the impact of cost-effectiveness analyses on their health plan coverage.

If IfPA can provide further feedback or information regarding these comments, please contact us at 202-499-4114.

Sincerely,

A handwritten signature in black ink, appearing to read "Brian Kennedy".

Brian Kennedy
Executive Director

Date: January 17, 2018

RE: Innovation and Value Initiative (IVI) Rheumatoid Arthritis (RA) Model - Response to Request for PublicComments

The following information is provided in response to request for public comment and is not intended as an endorsement of any usage not contained in the Prescribing Information. For complete information, please refer to the full Prescribing Information for each product, including the following sections: BOXED WARNING(S), INDICATIONS AND USAGE, DOSAGE AND ADMINISTRATION, CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS, and ADVERSE REACTIONS.

Contact Information

Submitter	Meredith Abraham, PharmD
Profession	Associate Director, Medical Information
Organization	Janssen Scientific Affairs, LLC
City	Horsham
State	Pennsylvania
Phone Number	215.325.2289
Email Address	Mabraha2@its.jnj.com

GeneralComments

The model contains a critical error because it does not consider SIMPONI ARIA as a treatment option. While the model does consider SIMPONI subcutaneous (SC), it is important that the developers of the IVI model recognize that SIMPONI SC and SIMPONI ARIA are entirely different formulations each with different pharmacokinetic, efficacy, and safety profiles, separate Phase 3 trial programs, and with separate US labeling as determined by the FDA. These products and their respective data sets must be considered separately (See attached SIMPONI Prescribing Information, SIMPONI ARIA Prescribing Information).

According to the report, comprehensive options to test different structures, data and assumptions are included in the model. However, in the online interface, the users do not have access to the options described in the report, and it is not clear what structure, assumptions and data sources are used in the online interface model. The danger in this is that the online interface may invite exploration without context or full understanding of how variables in the model relate which may cloud interpretation of results. For example, model assumption H2 (Stephens et al, 2015) uses methodology that is highly specific to patients with early aggressive RA, a population that was not tested in trials of golimumab or infliximab and most other biologics. This may be why significant variation in change in HAQ progression is noted with the H2 model assumptions vs H1 and H3.

In general the model in its current state may be too complicated for non-expert users. Language should be added to the online version to caution use by those unfamiliar with modelling methodologies.

Overview of IVI RA Model

The IVI-RA model is not a value assessment framework but a model that simulates the costs, health outcomes, and risks associated with treatments for RA. It can therefore be used with any value framework preferred by the user. The online tools support both cost-effectiveness analysis (CEA) and multi-criteria decision-analysis (MCDA).

Value Assessment

In MCDA, number of serious infections is included as one criteria. However, serious infections are also included in the CE model and have cost and quality of life impact already. The impact of serious infection in the value assessment appears to be counted twice. It should be removed from either CE model or the MCDA criteria to avoid double counting.

Populations

Page 18, Paragraph 2- Data from the CORRONA registry may underestimate the disease burden in the US RA population related to disease activity and physical function. Note Table 2b in Curtis et al which shows lower HAQ and CDAI scores compared other data sources. We recommend not depending on a single source in order to avoid outliers and to identify mid-range estimates for population disease activity (Curtis et al, 2010).

Source Data and Parameter Estimation

1. NMA: The inclusion and exclusion criteria are not described. Trials included in the NMA differ from other published studies such as by the Institute for Clinical and Economic Review (ICER) and the results are also considerably different. As stated previously, SIMPONI ARIA and SIMPONI SC must be considered as distinctly separate products. Missing GO-FURTHER trial data for SIMPONI ARIA are provided in Table 1 below and should be included in the NMA. Differences in patient characteristics and trial design differences are not considered in the NMA. For example, in the RAPID 1&2 trials, patients who did not achieve ACR 20 at weeks 12 and 14 were designated treatment failures and were withdrawn from the study at week 16. Other trials including GO-FURTHER called for treatment failures to continue in the trial. This leads to considerably low placebo rates at week 24 in RAPID 1&2 trials compared to other trials. As NMA did not adjust for this difference, the results are inevitably biased in favor of Cimzia (see reports published by the Canadian Agency for Drugs and Technologies in Health cited in References below). In further proof of the inflation of clinical benefit of RAPID 1&2, the head to head comparison of Cimzia vs Humira (EXXELERATE trial) did not meet its primary endpoint for superiority, and demonstrated no statistically significant difference in efficacy between Cimzia and Humira in combination with MTX in both short-term (12-week) and long-term (2-year) evaluations (Smolen et al, 2016).
2. The model considers dose adjustment for REMICADE but not for Humira or Actemra although labels for each of these products allow for dose adjustment. Dose adjustment can make a material change in the cost attributed to each therapy. Input options for dosing should be allowed from two perspectives: the average drug quantity used per administration and the average dosing frequency (or number of doses per year) during maintenance therapy. These input options to the model will allow users to understand the impact of dose escalation using their own existing utilization data.

The following biologics have dose escalation options included within their product labels (See respective product Prescribing Information):

Humira: Available references have found dose escalation of Humira to occur in 12.6-24.3% of patients (Fisher et al, 2013).

REMICADE: REMICADE provides dosing flexibility for patients with RA. The REMICADE PI recommends that patients be initiated on 3 mg per kg with maintenance dosing every 8 weeks and allows dose adjustment to as high as 10 mg per kg or maintenance dosing frequency as often as every 4 weeks. In real world practice the average dose of REMICADE has been reported to be 5.5 mg per kg. See Bolge et al for dose escalation assumptions that may be more consistent with clinical practice over the course of a 12-month period (REMICADE Prescribing Information; Bolge et al, 2012).

Actemra IV Administration: When used in combination with DMARDs or as monotherapy the recommended starting dose is 4 mg per kg every 4 weeks followed by an increase to 8 mg per kg every 4 weeks based on clinical response (Actemra Prescribing Information).

P 30, 8.4 - Rate of HAQ Progression for cDMARDs vs tDMARDs:

8.4.1 : HAQ progression parameters of cDMARD-treated patients are derived from Wolfe and Michaud 2010 and are "assumed to reflect the course of progression of HAQ in the absence of tDMARDs". Our reviewers suggest that HAQ trajectory of cDMARD patients before progression to tDMARDs does not adequately represent the presumed trajectory of cDMARD patients' HAQ in the absence of tDMARDs availability. While this may be a minor issue if this estimator is used in a model run intended to derive relative cost effectiveness of differing treatment sequences including line 2 biologics, this is a minor issue it becomes more significant when the model user is interested in ascertaining cost effectiveness of tDMARDs overall, as compared to cDMARDs. In this scenario, the model, and this estimator, will result in significant under-valuing of tDMARDs as this estimator will not capture true HAQ trajectory of patients in an environment of only cDMARD use.

8.4.2 : HAQ progression parameters of cDMARD patients are alternately used from Norton et al 2013, 2014. The HAQ trajectory of 1 of 4 cohorts cited in this paper represent the HAQ trajectories of patients with lower disease severity, and which are partially on biologics. As with issues in 8.4.1: Use of this parameter is an imperfect match for the decision analytic question that this model seeks to address. Similar to above comments: If the model user intends to compare relative cost effectiveness of two biologics after a presumed 1st course of cDMARDs (as a treatment sequence comparison), then this estimator mismatch may not present as significantly biased to cost effectiveness between tDMARDs. If, however, the user is interested in the question of overall cost effectiveness of tDMARDs compared to cDMARDs, then this parameter significantly underestimates the presumed HAQ trajectory of a general population on cDMARDs who have not accessed tDMARDs. This will significantly under-value the utility of tDMARDs.

The validity of this model to evaluate overall tDMARD cost effectiveness versus cDMARD is seriously compromised by the lack of a HAQ trajectory parameter than truly measures expected HAQ trajectory of cDMARD patients in the absence of tDMARD availability. This key parameter may be derived from data that may be available for HAQ trajectories before tDMARD introduction (eg from 80s or 90s), or from recent patient data from countries where patients may not have had access to tDMARD treatment, but for whom the imperative of aggressive early treatment via cDMARDs may have been presumably understood.

P 41 – HRU Costs are derived from citations in the literature and are inflated to 2016 values. This may lead to inaccuracies since literature citations used may be several years old. It would be preferred if inputs for more recent data could be obtained from contemporaneous databases or input by the user to minimize error introduced by inflation assumptions.

P 33 – Treatment discontinuation derived from CORRONA is an average of all anti-TNF agents, yet it is well documented that treatment discontinuation rates are higher for subcutaneously administered anti-TNF agents as compared to REMICADE. In Greenberg et al 2012, treatment persistency for bionative patients who initiated REMICADE was 76% at 12 months and 63% at 24 months, compared to 68% and 53% for Humira at 12 and 24 months, respectively and 72% and 53% for Enbrel at 12 and 24 months, respectively. Additional sources for treatment persistency include Fisher et al 2013 and Chastek et al 2016 (references provided below). Furthermore, we have shown that discontinuation rates for SIMPONI ARIA are similar to that of REMICADE (Data on File). Persistency for SIMPONI SC was cited as 75.3% in a study by Mourão et al, 2016.

P 42, Table 16 - Wholesale Acquisition Costs (WAC) are not current for some biologics. Drugs, effective dates and WAC are noted below.

- a. Enbrel- 1/1/18 \$1,218 per 50 mg syringe; Humira 1/1/18 \$2,436.02 per 40 mg syringe; REMICADE 2/9/17 \$1,167.82 per 100 mg vial; SIMPONI SC - 1/3/18 \$4,519.76 per 50 mg injector; SIMPONI ARIA (intravenous) 9/6/17 \$1,652.59 per 50 mg vial; Cimzia 1/1/18 \$4,044.32 per 400 mg; Actemra IV- 1/1/18 \$1,042.00 per 200 mg vial; Actemra SC - 1/1/18 \$984.72 per 162 mg syringe; Orencia IV 1/1/18 \$1,046.25 per 250 mg vial; Orencia SC - 1/1/18 \$1,032.76 per 125 mg syringe; Rituxan 1/1/2018 \$903.38 per 100 mg vial; XeljanzXR 1/1/18 \$4,095.64 per 30 d supply of 11 mg tablets.
- b. Drugs provided through a medical benefit or Medicare Part B are typically reimbursed using Average Sales Price (ASP). The largest segment of Janssen RA products, (REMICADE and SIMPONI ARIA), are most commonly covered under a medical benefit. The ASP calculation captures the list prices and the non-statutory discounts provided for all manufacturers. ASP is published quarterly and is available on the CMS website, but lags in time by approximately 2 quarters.
- c. In Table 16 (P. 42) dose escalation is assumed for REMICADE but not for Humira although Humira PI does allow for dose escalation in patients not taking MTX (Humira Prescribing Information). We recommend that the base case dosing assumptions conform exactly to the label and that the model allow the end user to vary the dosing and dosing interval assumptions rather than force the end user to compare to pre-set dosing and administration assumptions that may not be reflective of their true utilization.
- d. Dosing for SIMPONI ARIA should be included:
 - i. The PI for SIMPONI ARIA specifies 2 mg per kg at weeks 0, 4 and every 8 weeks thereafter (SIMPONI ARIA Prescribing Information).
- e. Dosing for administration of IV Actemra should be included:
 - i. IV Actemra dosing per the PI is 4 mg per kg every 4 weeks followed by an increase to 8 mg per kg every 4 weeks based upon clinical response (Actemra Prescribing Information).

Limitations and Areas for Improvement

P 8 - The model utilizes serious infection rates and changes in HAQ score during the first 6 months from baseline based on clinical trial evidence. It is known that differences in clinical trial populations and adverse event rates differ in those biologics that were first to market vs those that entered later. This could be a confounder in the model.

P 8 - The model assumes patients change treatments if they have a serious infection. This may not be consistent with real world scenarios where treatment might be held and then restarted.

P 9 - The Wailoo regression algorithm may have limitations for HAQ mapping. Pennington et al reported that the use of this utility mapping algorithm may have impact on overall results, particularly on inflation of the magnitude of the ICER (Pennington et al, 2013).

Appendices

P 67, Table A9- Add missing ATTRACT 102 week publication:

- Maini RN, Breedveld FC, Kalden JR, et al. Sustained improvement over two years in physical function, structural damage, and signs and symptoms among patients with

rheumatoid arthritis treated with infliximab and methotrexate. *Arthritis Rheum.*
2004; 50: 1051-1065.

P 70, Table A10- The correct n for the REMICADE 3 mg/kg every 8 weeks +MTX treatment arm is 86.

P 70, Table A10 - Add missing ACR20 and HAQ-DI data from ATTRACT at week 20 (Data on File, provided in Table 1)

P 71, Table A10 - The correct n for the SIMPONI 50 mg +MTX treatment arm in GO FORWARD is 89.

P 71, Table A10 - Add missing DAS28 data at week 24 from GO FORWARD (Data on File, provided in Table 1)

P 71, Table A10 - Add missing ACR20, ACR70, DAS28, and HAQ-DI data at week 24 from the GO FURTHER trial (Data on File, provided in Table 1)

Table 1: Missing or Corrected Efficacy Data Related to Janssen Products

Trial ID	Treatment	N	Time in Weeks	ACR20 n(%)	ACR 50 n(%)	ACR 70 n(%)	ΔDAS28 (SE)	ΔHAQ-DI (SE)
ATTRACT	REMICADE 3 mg/kg Q8W+MTX	86	30	43(50%)	22 (27%)	7 (8%)	Not available	Not available
	MTX	88	30	18(20%)	4 (5%)	0 (0%)	Not available	Not available
GO FORWARD	SIMPONI 50 mg+MTX	89	24	79 (60%)	49(37%)	27(20%)	DAS28 CRP- Median change from baseline (IQR): -1.682 (-2.539, -0.676) Mean change from baseline (SD): -1.621(1.2873)	-0.38 (0.16)
	MTX	89	24	25(28%)	12(14%)	5(5%)	DAS28 CRP- Median change from baseline (IQR): -0.624 (-1.469, 0.215) Mean change from baseline (SD): -0.692(1.3206)	-0.13 (0.13)
GO FURTHER	SIMPONI ARIA 2 mg/kg +MTX	395	24	248(63%)	138(35%)	69(18%)	DAS 28 CRP- Median change from baseline (IQR): -2.0441 (-2.9928,	HAQ-DI Median change from baseline (IQR): 0.50 (0.1250, 0.8750)

REFERENCES

Actemra® (tocilizumab) Prescribing Information.

Bolge SC, Carter CT, Lofland J, et al. Comparative multidatabase analysis of dosing patterns and infusion intervals for the first 12 infliximab infusions in patients with rheumatoid arthritis. *Clin Ther*. 2012; 34(12):2286-2292

Canadian Agency for Drugs and Technologies in Health. *Therapeutic Review: Biologic Response Modifier Agents for Adults with Rheumatoid Arthritis*. July 2010. Available at:
https://www.cadth.ca/sites/default/files/pdf/TR_RA_Clinical_and_Economic_Overview_e.pdf

Canadian Agency for Drugs and Technologies in Health. *Common Drug Review: Certolizumab Pegol (Cimzia-UCB Canada Inc)*. May 2010. Available at:
https://www.cadth.ca/media/cdr/complete/cdr_complete_Cimzia_May-28-2010.pdf

Chastek B, White J, Van Voorhis D, et al. A Retrospective Cohort Study Comparing Utilization and Costs of Biologic Therapies and JAK Inhibitor Therapy Across Four Common Inflammatory Indications in Adult US Managed Care Patients. *Adv Ther*. 2016;33:626–642.

Curtis, JR, Jain, A, Askling, J, et al. A comparison of patient characteristics and outcomes in selected european and us rheumatoid arthritis registries. *Seminars in Arthritis and Rheumatism*. 2010 (40):2-14.

Data on File. Janssen Biotech, Inc.

Fisher MD, Watson C, Fox KM, et al. Dosing patterns of three tumor necrosis factor blockers among patients with rheumatoid arthritis in a large United States managed care population. *Curr Med Res Opin*. 2013;29(5): 561–568.

Greenberg JD, Reed G, Decktor D, et al. A comparative effectiveness study of adalimumab, etanercept and infliximab in biologically naïve and switched rheumatoid arthritis patients: results from the US CORONA registry. *Ann Rheum Dis*. 2012;7: 1134-1142.

Humira® (adalimumab) Prescribing Information.

Keystone EC, Genovese MC, Klareskog L, et al. Golimumab, a human antibody to tumour necrosis factor alpha given by monthly subcutaneous injections, in active rheumatoid arthritis despite methotrexate therapy: the GO-FORWARD study. *Ann Rheum Dis*. 2009;68: 789-796.

Maini R, St. Clair, EW, Breedveld F, et al. Infliximab (chimeric anti-tumor necrosis factor alpha monoclonal antibody) versus placebo in rheumatoid arthritis patients receiving concomitant methotrexate: a random phase II trial. *Lancet*. 1999;354: 1932-1939.

Maini RN, Breedveld FC, Kalden JR, et al. Sustained improvement over two years in physical function, structural damage, and signs and symptoms among patients with rheumatoid arthritis treated with infliximab and methotrexate. *Arthritis Rheum*. 2004; 50: 1051-1065.

Mourão AF, Ribeiro C, Borges J et al. Real-life effectiveness of golimumab in biologic-naïve rheumatoid arthritis patients - data from Reuma.pt, a Portuguese registry. AbstractAB0297 published at the Annual European Congress of Rheumatology (EULAR 2016).

Pennington B, et al. The influence of HAQ utility mapping algorithms on the cost-effectiveness of second line biologics for treatment of rheumatoid arthritis. *Value in Health*. 2013:A323–A636.

[REMICADE® \(infliximab\) Prescribing Information.](#)

[SIMPONI ARIA® \(golimumab\) Prescribing Information.](#)

[SIMPONI® \(golimumab\) Prescribing Information.](#)

Smolen JS, Burmester G, Combe B, et al. Head-to-head comparison of certolizumab pegol versus adalimumab in rheumatoid arthritis: 2-year efficacy and safety results from the randomised EXXELERATE study. *Lancet*. 2016;388:2763-2774.

Weinblatt ME, Bingham III, CO, Mendelsohn AM. Intravenous golimumab is effective in patients with active rheumatoid arthritis despite methotrexate therapy with responses as early as week 2: results of the phase 3, randomised, multicentre, double-blind, placebo-controlled GO-FURTHER trial. *Ann Rheum Dis*. 2013;72:381–389.



Pfizer, Inc.

Received via email from Sachin Kamal-Bahl, PhD

As a strategic advisory board partner we are grateful for the opportunity to provide additional comments on the on the IVI RA model. We continue to see a unique opportunity in your approach to create an open source model for all to have access to. In reviewing the documents provided we present the following comments. We continue to suggest that IVI take the time to address these comments, review the changes made with the advisory board including all stakeholders, and integrate a formal process with approvals from stakeholders as you continue to advance the model forward.

For clarity we have separated the comments into sections 1) general comments 2) model specific comments.

General comments

- 1) We ask that IVI continue to review the principles around value frameworks that have been developed by PhRMA and NPC. These principles should form the underlying basis of both how IVI approaches model development, and what key parameters IVI seeks to advance in its models. Currently, we see significant opportunity for IVI to improve on its alignment against these principles. For example: a key consideration in both PhRMA and NPC's principles revolves around transparency in process. ICER et al largely build models behind closed doors, without gathering feedback from stakeholders as the inputs are selected. When they do release their draft findings, stakeholders are left struggling to understand complex models in often short review periods. Unfortunately, IVI has made the same errors in its process. Although additional data has been provided in this updated version, the sourcing and justification of decisions remain unclear in many instances which reduces the ability to understand or comment on the appropriateness of the specific parameters. We reiterate a strong believe in open source not only in the development of the model structure but also in the parameters used to populate the model.
- 2) Although, many front end descriptions have been developed the many front end tools can create confusion in the users' ability to adapt or amend data. Of significant note, depending on the front end used, there can be a lack of information on the efficacy data, data used in the model parameters, and an overall lack of clarity as the application of the correct technical documents all of which make using the model online dashboard a confusing process. Depending on the front end used there are significant differences in the changes that can be made and as such further clarity should be provided to the end user.
- 3) We applaud the integration of multi-criteria decision analysis and the ability of the end user to adjust preferences weights. This is a strong addition to the tool as is specific domains as preference for treatment administration, insurance, and time since approval. We would suggest that an additional function should be the depth of indications each product has. Given that a medication which could be used in mulitpile comorbid conditions, the insurance value of the product could be improved.

- 4) We are happy to see greater integration of patient level information. However, while the model is recommended to be built on consensus, information on the consensus process, beyond a description of the use of patient level input, has not been provided. As noted above, we continue to suggest that IVI take the time to address these comments, review the changes made with the advisory board including all stakeholders, and integrate a formal process with approvals from stakeholders as you continue to advance the model forward.
- 5) As previously suggested, we would also suggest that a number of external rheumatology focused health economists should be integrated as an additional advisory board to IVI to continue to advance this model. We would be happy to submit the names of number of individuals who could provide independent, objective feedback and support to strengthen the project. This would further separate this work from past efforts.
- 6) IVI's objective was to develop models that were market driven; however the model as it is developed does not appear to reflect the US environment and does not seem to integrate such aspects as payer restrictions, step edits, etc. An innovative model should allow for an assessment of these processes and the impact of removing these restrictions on the identified endpoints of interest to patients, rheumatologists, and relevant stakeholders
- 7) Further, given access to the above data, one should use it and only resort to modeled results when measured outcomes are not present. This would make the result far more valuable to all interested stakeholders and stand out from past efforts to this end.

Model specific comments

1) Model Data used

Given the maturity of the data on many advanced RA therapies including advanced therapies such as Anti-TNFs, novel Mechanism of Action Biologics, and JAK inhibitors, we would suggest that a truly innovative approach should include a partnership with the developer of an established observational data set to better achieve the outlined goals of a transparent, market driven, multi-perspective model. Similarly, we recommend that data be requested from stakeholders (industry, payers, and registry owners) which also could inform the model(s) being developed. Regardless, it would be preferable to integrate a real world data set/registry which could report on the changes of disease activity and likelihood of switch to better reflect the treatment patterns in the US.

2) Model structure

- a. The switching methodology and justification for treatment switching is poorly outlined and does not reflect current clinical practice or how physicians make the decision to change therapy in the US healthcare system. Most notably it ignores the treat to target ideology as recommended by the American College of Rheumatology.
- b. The report refers to the 336 possible model structures, however many of these structural changes are based on the estimation of how HAQ is derived (either by ACR response, or EULAR response) or treatment response is determined (ACR,

DAS28, SDAI, or CDAI). This appears to be an attempt to integrate the methods of efficacy of a number of other published CEA models rather than building a consensus on one type of structure/modelling method. Also, as outlined further the transformation from ACR to other variables of effectiveness is flawed.

- c. The issue of taking a categorical variable ACR 20/50/70 response to a continuous variable such as SDAI, CDAI, DAS28, and HAQ is not clearly outlined or described in the document. Algorithms for mapping the ACR response to SDAI, CDAI, DAS28, EULAR response, and HAQ are taken from a number of different, small mapping trials which may or may not show valid relationships between the measures. This limitation was discussed in detail in the document, but further information and variability in the estimates (and the distributions which could be used in the PsA) should be included in the description and model.

3) Patient population

- a. The initial data on the patient population were taken from Curtis et al and US301 Clinical trial. Of note is that is unclear which of the US registries reported on in Curtis et al is being used and it appears that difference data points (age, HAQ score, etc.) are taken from different registries without a single registry's data being identified as the seminal piece of patient characteristics. Similarly, reviewing the data from Curtis et al it is unclear of when characteristics were collected- at the time of treatment initiation or registry recruitment. A more appropriate source of data would be a real world registry which would report on the clinical characteristics of a patient population starting an advanced therapy. These data should be easily accessible through collaboration with a US registry such as Corrona.
- b. It also questionable to take the data from a leflunomide (a csDMARD) trial completed published 14 years ago- US301 -to inform a model designed to report on advance therapy utilization. This trial and trial data may not reflect the current patient population or treatment patterns based on disease activity. Furthermore, the work completed by Smolen et al only for reported on the validity of SDAI not CDAI.
- c. The algorithm for distributions used in the heterogeneous patient population also could have been replaced simply by using a patient population taken from US based registries who are initiating an advanced therapy. Similarly, the use of a RWD source of advanced therapy experienced (referred to as a bDMARD experienced) patients could inform on the baseline characteristics for those patients when entering the model.
- d. It is also noted that disease duration is included in baseline characteristics of RCTs and is integrated in CEA models. The justification of omission of this variable in the patient populations should be better articulated. Given that disease duration is identified in the latent class growth model for HAQ progression as a key variable it is unclear how that has been integrated into the model.

4) Comparators

- a. We question the use of csDMARDs as a comparator for anything other than salvage or last line of therapy. A patient whose disease would necessitate an advanced therapy is likely to receive that therapy and while economic theory does propose a “do nothing” strategy we feel this is both an unethical and unlikely strategy and does not reflect clinical practice. Engagement with the recommended panel of RA specialists would help clarify this point.
- b. As described in the general comments, the structure of the model does not appropriately describe the system in the United States for an RA patient. The current formulary and PBM processes which often include preferential products which impact available treatments. A more appropriate model would include data reflecting that and changing the process to assess the impact of these step edits and restrictions of products.
- c. It is also noteworthy that although many treatments for RA have the opportunity to increase dose or decrease the time between doses, this is not included in the modeling. Again, a number of studies using evidence from registries/observational data has reported on this topic and could be integrated into the model.

5) Efficacy data

- a. Initial treatment response, within the model for the initial 6 months of any treatment the estimated response was taken from a newly completed IVI Network Meta Analysis (NMA) using only using RCT data. However, very little information on the NMA is available, of specific note is the background information on the methods of study selection, the studies and data included in the NMA, and methods employed to run the NMA. It should be recognized that a number of registry, non-RCT, and observational trial data sources exist which include US and non-US populations, however, based on the limited information provided it appears that these data have been excluded from these NMA and therefore are ignored. This appears to go against the ideology of reflecting the current US market/being market driven.
- b. For the initial treatment phase it is further unclear why only RCT data were used when other components of the model integrate observational data (i.e. Adverse Event data, initial populations, relationship between HAQ and costs of RA). Given the long history of data and the integration of other registry data in other aspects of the model (i.e. Adverse events), it is unclear why a NMA was required and one so restrictive of inclusion criteria.
- c. Similarly, it is unclear why a factor was applied for the reduction of initial treatment response when patients in the model are assumed to have previous biologic or advanced therapy exposure (either because they have switched treatments in the model or entered the model with the baseline characteristic of having previous advanced therapy or biologic exposure). A number of RCT studies and observational data sources exist to more accurately estimate the different efficacy for a patient who has been exposed to previous biologic or advanced therapy vs a patient who is naïve to these treatments. These data are

ignored by this model (or else perhaps have been inappropriately integrated into the NMA).

- d. In the description of efficacy and progression of disease in the period after the first 6 months the method described a number of sources of data. Many of these described methods have been used in previous cost effectiveness models, but appear to ignore the progression of the disease and the natural path of RA observed in clinical practice.
- e. For the data on the HAQ progression (in absence of or in the presence of bDMARD or tsDMARD) these data are taken from a number of different registries and sources of data nationally and internationally. Again, a partnership with a US registry or source of real world data would better articulate the disease progression, as well as provide greater information on the thresholds of change for switching treatments.

6) Adverse event reporting

- a. Using a single adverse event (serious infections) does not represent the many events, many of which may require healthcare resource utilizations, and may or may not require discontinuation of treatment. A number of trials (observational and RCTs) and registries have reported on the safety of the treatments included in the model. Using a single adverse event, having it cause a change in treatment, while this is stated as a model limitation, this does not reflect clinical practice. Furthermore this ignores a number of studies which have been completed in RA patients receiving advanced therapies reporting on the risk and changes in risk reported by treatment for a number of unique adverse events and co-morbidities.
- b. Additionally, it is assumed that the risk of adverse event is identical for all treatments. This also ignores a significant amount of literature on the safety of advanced therapies.

7) Utility measurement

- a. The model reports on two models for transforming HAQ score to utility. It is noted that some other methods were ignored due to the data used coming from clinical trial data. This exclusion of methods seems peculiar since clinical trial data is identified as being the primary tool for efficacy measurement. Similarly, studies measuring the impact of using different utility measures (SF-6D, HUI2/3, etc.) have been completed but were ignored in this model. Similarly a number of RCTs and registries have collected and reported on utility data which could have been integrated into this modeling exercise.

8) HAQ to mortality and healthcare resource utilization

- a. Mortality, RA-related hospitalizations, and productivity costs are all estimated from the patient HAQ score. While this has been used in a number of previous CEA models, again, there is potential to use data from an established registry or data set to continue to improve the reporting of this topic.

9) Prices and costs

- a. Currently the model uses the WAC price which we feel is not appropriate. The WAC, as published by FDB represents the manufacturer's (for purposes of this Drug Price Policy, the term "manufacturer" includes manufacturers, repackagers, private labelers and other suppliers) published catalog or list price for a drug product to wholesalers as reported to First Databank by the manufacturer. WAC does not represent actual transaction prices and does not include prompt pay or other discounts, rebates or reductions in price. First Databank does not perform any independent investigation or analysis of actual transaction prices for purposes of reporting WAC. First Databank relies on manufacturers to report or otherwise make available the values for the WAC data field.

An alternative source of data would be the NADAC; it is our position that the NADAC is a more appropriate measure of drug price. The purpose of the NADAC is to create a new national price benchmark that is more reflective of the prices that pharmacies pay to acquire prescription and over-the-counter drugs. The statute provides that such prices represent a nationwide average of consumer purchase prices, net of discounts and rebates. The survey data will provide information which CMS expects to use to assure compliance with Federal requirements. A monthly nationwide survey of licensed retail community pharmacies, which will include independent pharmacies and chain pharmacies in the United States, will be performed to collect drug acquisition cost information. To ensure that NADACs are accurate, timely, and robust, the NADACs will be reviewed and updated on a weekly basis.

- b. The integrated use of only productivity as the societal measure of cost may not fully articulate the costs of absenteeism, presenteeism, and other non-monetary or intangible costs. Further articulation of the costs used would assist a proper assessment of the model.

Darius Lakdawalla, PhD
Executive Director
Innovation and Value Initiative
11100 Santa Monica Blvd, Ste 500
Los Angeles, CA 90025

January 17, 2018

RE: Innovation and Value Initiative (IVI) Open-Source Value Project: IVI-Rheumatoid Arthritis (RA) Model Request for Public Comment

Dear Dr. Lakdawalla,

Sanofi-Genzyme and Regeneron appreciates IVI's efforts to solicit and integrate stakeholder feedback into the IVI-RA Model, the first project released through the IVI Open-Source Value Project. We are pleased to see flexibility of the model, in which many inputs and assumptions can be adapted by the user, and that the model is detailed in the technical document entitled, "*A Description of the IVI-RA Model v1.0.*" We also commend the development team for creating multiple ways to use the IVI-RA model based on level of expertise, type of user or stakeholder, and intended use of the information. The IVI-RA Value Tool, IVI-RA Model Interface, and the iviRA R Package enhances the transparency of the Open-Source Value Project.

We recognize the extensive effort and investment of resources it took to undertake this project in a disease state such as rheumatoid arthritis. With 384 available model structures, many scenarios of interest to the end user will be accounted for by this model. Based on our review of the IVI-RA Model, our comments and suggestions below focus on the following: *model functionality, user navigation, IVI-RA Value Tool population, treatment sequences, model structure, parameter values, cost-effectiveness analysis and future adaptations.*

Model functionality

The model interface allows the user to modify input settings in the Model Setup section. It would be helpful to have functionality built-in to allow the user the option to save the modified inputs entered into the model and to have access to the saved data upon accessing the model again or after running an alternative scenario. Similarly it would be helpful to be able to save outputs of the model results.

User navigation

The "Introduction" tab indicates that the user can modify the simulation settings in the "Setup Model" tab, however "Simulation Settings" is found in the "Run Simulation" tab. We recommend clarifying this under Custom analysis in the "Introduction" tab. Clarity is needed to inform the user where "modify output settings" can be found. The "Introduction" tab under "View Model Results" in "Custom analysis" section implies that the user would click on "Modify output settings" to select between societal or health care sector), however it appears that the selection is made in "Value to the Healthy" and "Cost-effectiveness Analysis" under the "View Model Results" tab.

We recommend providing a statement in the Model Interface “Introduction” tab regarding where the user can find references for the default settings (input parameters), as well as alerting the user that parameters will be referenced on the “View Inputs Used in Simulation” tab. Ideally, each default parameter should be referenced on each page of the “Setup Model” section.

IVI-RA Value Tool population

Within the Value Tool, concepts are clearly defined so that the user understands the inputs, outcomes, value calculations and scenario analyses. In the “Setup” tab of the Value Tool, it would be helpful to also describe the default population simulated in the tool for those decision-makers who do not intend to make adjustments to the parameters on that tab. This will assist the user with a better understand of what is being reviewed when “Pick for me” is selected and assist with appropriate utilization of the data.

Treatment sequences

KEVZARA® (sarilumab) was approved by the FDA in May 2017 for the treatment of adult patients with moderately to severely active rheumatoid arthritis (RA) who have had an inadequate response or intolerance to one or more disease-modifying antirheumatic drugs (DMARDs). We therefore recommend consideration of its inclusion in the treatment sequence of the model. For more information, please refer to the KEVZARA full prescribing information, including **Boxed WARNING**. We would also like to recommend the user have the capability to add new medications, dosage forms and frequency to the model in the event approval is granted prior to model revision.

Model structure

While the IVI-RA Model is very flexible in the model structure, we would like to recommend that users have multiple options for utility equations, in addition to the Hernandez-Alava mixture model and Wailoo logistic regression equation. Other options for consideration can be found in cost-effectiveness analyses such as Bansback et al. 2005, Hurst et al. 2006, Malottki et al. 2011, Brennan et al. 2004, Ducournau et al. 2009, and Boggs et al. 2002 (Table 1). Similarly, we recommend the flexibility of modifying mortality rates based on the impact of baseline functional status on mortality, as suggested by Yelin et al. 2002, Michaud et al. 2012, Diamantopoulos et al. 2012, or Marra et al. 2007.

Table 1. Examples of Utility Equations*

	BRAM	Abbott	Wyeth Pharmaceuticals	Schering-Plough Ltd	Roche	Bristol-Myers Squibb Ltd
Mapping of effectiveness data to utility	Quadratic equation using dataset supplied by Hurst and reported in Hurst et al. 1997 in the absence of any more recent dataset available to the assessment group	HAQ scores were converted to EQ-5D scores according to equations (linear and non-linear) developed by Ducournau et al. 2009 using data from TOC trials. The non-linear equation was used for the base-case analysis, while the linear equation was examined in sensitivity analyses	HAQ scores were converted to EQ-5D scores according to a linear equation developed by Brennan et al. 2004	Utility was estimated to be a function of EULAR response, treatment (on biologic treatment or not), health-state utility at time of treatment initiation, age, disease duration, number of previous DMARDs and gender	HAQ scores were converted to EQ-5D scores according to the non-linear equation developed by Ducournau et al. 2009 using data from TOC trials	HUI 3 utilities were calculated from the HAQ based on a conference abstract [Boggs 2002] EQ-5D utilities calculated from HAQ were used in a sensitivity analysis

*Source: Table 74 Data input and assumptions used in manufacturer models
Malottki et al. Health Technol Assess 2011;15(14)

Parameter values

The user is allowed to adjust multiple parameters within the Treatment Cost section of the model interface, including accounting for a discount (lower and upper). An additional functionality of accounting for patient copay may be appreciated by some stakeholders/decision-makers using the model interface as well as in the Value Tool.

While users are allowed to adjust parameters for time to treatment discontinuation, we recommend allowing for customizable treatment discontinuation event rates for each medication included in the model. This provides greater flexibility for the user to incorporate real world data reflective of their organization.

Cost-effectiveness analysis

The model results section provides an extensive number of results allowing potential users a variety of data to select and utilize based on their organization's specific needs to inform decisions. Within the cost-effectiveness analysis section, some users may be interested in viewing individual medication incremental cost-effectiveness ratios (ICERs) rather than seeing the ICER of sequences in order to inform decisions. We recommend the incorporation of individual ICERs in addition to sequence ICERs.

Model Interface

In the Model Interface, we are unclear about the adjustment of the initial treatment effect according to treatment line. Specifically, how are the low and high values of the adjustment factor applied in the model? Additionally, while the Model Interface is geared toward experienced modelers, there are some parameters such as the ACR and EULAR response rates for initial treatment responses that are not

available to the user. Instead, the user is given the parameter estimates of statistical models (such as the Bayesian models for ACR and EULAR response rates), and therefore must perform calculations to understand the response rates used in the model. We are unclear how the user can enter alternative response rates.


Future adaptations

We appreciate the thoughtful construction of the model which facilitates utilization by a broad set of stakeholders within the US healthcare system. While the US healthcare system is the focus of the IVI Open-Source Value Project, a model further adjusted to allow for ex-US adaptations in the future would be of interest.


In conclusion, we appreciate IVI's effort to incorporate stakeholder feedback as an important part of the Open-Source Value Project's iterative process using an ongoing cycle of public feedback, expert review, revision, and re-release, and its willingness to engage in such a manner for all models constructed through the Open-Source Value Project.

We look forward to further discussions and encourage IVI to continue its efforts to apply scientific principles to the study of value in medicine and create open-source tools that support decision-making and inform health policy.

Yours Sincerely,



Sheila M. Thomas
Senior Director
Global Health Economics & Value Assessment
Sanofi



Vera Mastey,
Executive Director
Health Economics and Outcomes Research
Regeneron

References

- Bansback NJ, Brennan A, Ghatnekar O. Cost effectiveness of adalimumab in the treatment of patients with moderate to severe rheumatoid arthritis in Sweden. *Ann Rheum Dis* 2005;64:995–1002.
- Boggs R, Sengupta N, Ashraf T. Estimating health utility from a physical function assessment in rheumatoid arthritis patients treated with adalimumab. ISPOR meeting presentation, Rotterdam, the Netherlands, 2002.
- Brennan A, Bansback N, Reynolds A, Conway P. Modelling the cost-effectiveness of etanercept in adults with rheumatoid arthritis in the UK. *Rheumatology* 2004;43:62–72.
- Diamantopoulos A, Benucci M, Capri S, Berger W, Wintfeld N, Giuliani G, Ricciardi W., Economic evaluation of tocilizumab combination in the treatment of moderate-to-severe rheumatoid arthritis in Italy, *J Med Econ.* 2012;15(3):576-85.
- Ducournau P, Kielhorn A, Wintfeld N. Comparison of linear and non-linear utility mapping between HAQ and EQ-5D using pooled data from the tocilizumab trials OPTION and LITHE. Annual Meeting of the British Society of Rheumatology (BSR) Glasgow, 2009; poster 258.
- Hurst et al. in Chen, Y.F., Jobanputra, P., Barton, P., Jowett, S., Bryan, S., Clark, W. et al. A systematic review of the effectiveness of adalimumab, etanercept and infliximab for the treatment of rheumatoid arthritis in adults and an economic evaluation of their cost-effectiveness. *Health Technology Assessment* (Winchester, England) 2006; 10(42):iii-iv.
- Malottki K, Barton P, Tsourapas A, Uthman A, Liu Z. Adalimumab, etanercept, infliximab, rituximab and abatacept for the treatment of rheumatoid arthritis after the failure of a tumor necrosis factor inhibitor: a systematic review and economic evaluation. *Health Technol Assess* 2011;15(14).
- Marra CA, Marion SA, Guh DP, et al. Not all “quality-adjusted life years” are equal. *J Clin Epidemiol.* 2007; 60(6): 616-24.
- Michaud K, Vera-Llonch M, Oster G. Mortality Risk by Functional Status and Health-related Quality of Life in Patients with Rheumatoid Arthritis. *J Rheumatol* 2012;39;54-59.
- Yelin E, Trupin L, Wong B, Rush S. The impact of functional status and change in functional status on mortality over 18 years among persons with rheumatoid arthritis. *J Rheumatol* 2002;29;1851-1857.



Tufts University

Received via email from Peter Neumann and Josh Cohen

Hi Darius,

The IVI RA model is impressive. The range of “knobs” made available to the user to see how results change in response to alternative assumptions and outcome measures is fantastic. The interface is easy to follow, and the generated figures are great. Likewise, the technical documentation is very good.

I do have several relatively minor comments:

- While the simulation is running, it would be good to have a dynamic display indicating that something is happening. Conveying the progress towards completing the requested total number of iterations would be even better. When I first started playing with the model, I wasn't sure it was really working and started clicking on things before the simulation completed.
- The model allows the user to assign a negative dollar value to a QALY. I think the lower bound on value should be zero.
- One aspect of the navigation that wasn't clear to me was how to easily get back to the home page, or the page where the user can start the simulation. It should be easy for the user to “bail out” at any point and start all over again.
- It would be nice to be able to easily download generated figures and the underlying simulated values. That functionality may be there, but I may have missed it.
- While the technical documentation was very good in general, it could be a little clearer about what's going on with the PSA. Is each individual simulated multiple times with random draws for uncertain parameters? Does the simulation do anything to separately characterize heterogeneity (differences between individuals) and uncertainty (results that vary because of imprecise knowledge)? For example, it would be nice to know the median number of QALYs gained, and how uncertain the estimate of the median is.
- Finally – it's great, and absolutely crucial that the user has access to the source code. That means that a party that is concerned with RA (and well-resourced) can ultimately see exactly what the IVI model does. This model is very complex, however, and it would be good to provide at least some indication of how a more intermediate technical user can find his or her way around the model. For example, I am familiar with C-sharp and a number of other programming languages, but I couldn't find my way to the “top” of the model stored in GitHub. Admittedly, programming isn't my main expertise, but I feel that even someone at my level should be able to get around the model more easily. I am not saying that IVI needs to provide substantially more documentation. But something to get people started would go a long way.

-Josh

Appendix B. IVI-RA Model Technical Expert Panel Membership

The IVI-RA Model Technical Expert Panel featured perspectives from leaders in health economics, epidemiology, rheumatology, and patient communities.

Louis Garrison, PhD, Chair

Health Economist, Professor Emeritus, University of Washington

Dr. Garrison joined the School of Pharmacy faculty in 2004. For the first 13 years of his career, Dr. Garrison worked in non-profit health policy research first at the Battelle Human Affairs Research Centers (Seattle), and then at the Project HOPE Center for Health Affairs (Virginia), where he was the Director from 1989-1992. Following this, he worked as an economist in the pharmaceutical industry for 12 years. From 2002-2004, he was Vice President and Head of Health Economics & Strategic Pricing in Roche Pharmaceuticals, based in Basel, Switzerland. In October 2016, he became Senior Visiting Fellow at the Office of Health Economics, London, UK. Dr. Garrison's research interests include national and international health policy issues related to pharmacogenomics and personalized medicine, regulatory benefit-risk analysis, insurance, pricing, reimbursement and risk-sharing agreements, as well as the economic evaluation of pharmaceuticals, diagnostics, devices, surgical procedures, and vaccines, particularly as related to organ transplantation, renal disease, influenza, measles, obesity and cancer.

Stephen Mennemeyer, PhD

Health Economist, University of Alabama School of Public Health

Stephen T. Mennemeyer is a Professor at UAB in the Department of Health Care Organization and Policy in the School of Public Health and a Scholar in the Lister Hill Center for Health Policy. He is the Chief Science Officer for the Survey research Unit in the School of Public Health. Dr. Mennemeyer holds a Ph.D. in economics from the State University of New York at Buffalo. He has previously worked for Abt Associates Inc. as a senior economist and project director (1979-1989), for the Health Systems Agency of Western New York (1977-1979) as an economist, and for the School of Pharmacy at the State University of New York at Buffalo (1975-1977) as Assistant Professor. Since 2010 he has been a member of the Agency for Health Care Research and Quality, Healthcare System and Value Research (HSVR) study section.

Dr. Mennemeyer's work includes cost effectiveness or cost utility analyses regarding stroke rehabilitation, antiviral therapy for neonatal herpes simplex virus, and drug abuse treatment for the homeless as well as other studies on child obesity, health care labor, clinical laboratory competitive bidding and patient response to hospital mortality information. Recently he developed a discrete event simulation model to examine the cost effectiveness of earlier cataract surgery as a strategy for reducing motor vehicle collision. He is currently working on an Agent Based simulation model to examine how treating HIV + women for depression may affect their adherence to HIV therapy and the transmission of the HIV virus.

Kaleb Michaud, PhD

Epidemiologist, University of Nebraska Medical Center

Dr. Michaud serves as co-director of the National Data Bank for Rheumatic Diseases (NDB), the largest long-term open-cohort observational study of its kind with over 40,000 patients enrolled. Since 2007, he has been the principal investigator of the RAIN database composed of several rheumatology clinics in



the Rheumatoid Arthritis Investigator's Network, and has led the transition to using electronic tablets in the UNMC rheumatology clinic. He is co-founder and director of Clinical Informatics for the Arthritis Internet Registry (AIR), a partnership with the NDB, Arthritis Foundation, and Quest Diagnostics. As an active volunteer in the American College of Rheumatology and Association for Rheumatology Health Professionals, he has served 8 years in developing the Rheumatology Informatics System for Effectiveness (RISE) national registry. He also works closely with Dr. Ted Mikuls on the Veteran Affairs Rheumatoid Arthritis (VARA) registry. His primary research interests revolve around pharmacoepidemiology, health projects include: mortality in rheumatic diseases, cost-effectiveness of rheumatoid arthritis treatments, and impact of collecting patient reported outcomes via smartphones. He has received funding from the AHRQ, NIH, Arthritis Foundation, and Rheumatology Research Foundation, and is the director of the UNMC Rheumatology fellowship research program.

Elizabeth Perkins, MD

Rheumatologist at Rheumatology Care Center (RCC)

After finishing her medical degree at UTHSCSA, Dr. Perkins was accepted to the Medical College of Virginia at the Virginian Commonwealth University in Richmond to complete her residency in Internal Medicine. She specifically chose this residency program to complete a women's health track as part of her training. Ultimately, it was her experiences with tough autoimmune diseases that influenced her decision to subspecialize in rheumatology. As a natural problem solver, Dr. Perkins quickly gravitated to the complex diseases that her field presents. Rheumatology Care Center is at the center of Dr. Perkins's sustained vision to provide the best health care possible. Her practice continues to grow rapidly with the support of wonderful staff and medical community. Her leadership and advocacy spirit fuel her efforts in the American College of Rheumatology (ACR), Alabama Society of Rheumatic Diseases (ASRD), and Arthritis Foundation.

Carole Wiedmeyer, MBA

Rheumatoid Arthritis Patient Researcher

Ms. Wiedmeyer is an experienced researcher who has led and contributed to numerous studies in the transportation, technology, energy and health care industries. As a patient with rheumatoid arthritis, she has also recently contributed to several rheumatoid arthritis-related studies. Her research experience includes developing quantitative and qualitative research designs; developing research instruments and fielding plans; moderating focus group/small group discussions and individual interviews; conducting website, software, and app usability testing; supervising fielding and data processing; conducting analysis and report writing; and presenting research findings. She has extensive experience successfully fielding large studies in the U.S. and internationally using a variety of data collection methods.

In addition to her work as a patient researcher, Ms. Wiedmeyer serves as a Patient Governor with the Global Healthy Living Foundation's (GHLF) ArthritisPower initiative, which established a network of thousands of arthritis patients dedicated to participating in arthritis research. During that time, she spearheaded the formation of the GHLF Patient Partners in Research, which expands the pool of patient volunteers who are trained in research methods and available to collaborate directly with academic researchers on arthritis research.

Appendix C. Delphi Surveys

Delphi Survey #1

Introduction

The IVI-RA model is part of IVI's Open Source Value Project (OSVP), which is building an open, collaborative, and consensus-based process for the development of tools for value assessment. Models developed by the OSVP process are iterative, evolving as the science of value assessment advances and as new evidence becomes available. OSVP models are released and updated using a four-step process:

1. Public release of the model.
2. Invite feedback and suggested improvements to the model in a public comment period.
3. A panel of experts determines which of the evidence-based suggestions for improvement suggested in Step 2 should be implemented by means of peer-review and a formal voting process.
4. Revise the model based on the feedback from the technical expert panel in Step 3.

The first OSVP model evaluates treatments for rheumatoid arthritis.

Version 1.0 is IVI's first release of the IVI-RA model, and is designed to provide a starting point for open debate (Step 1). As such, the model is very flexible and allows users to choose from a large number of the plausible model structures supported by clinical practice and prior decision-analytic modeling research in rheumatoid arthritis.

Over the past few months we have received feedback and suggestions for improvements from different stakeholders (Step 2), which were discussed with you in a teleconference at the end of February. Based on the discussion we decided to proceed with a modified Delphi approach to get consensus which recommendations for model improvement should be implemented (Step 3). In this Delphi process we only focus on those suggestions for improvement that may require rather extensive resources to implement (such as elaborate programming or additional evidence collection) or are likely to have a substantial impact on the findings of a model analysis.

The modified Delphi process will proceed in the following way:

Survey 1:

- Each suggested change from the public comments were categorized by domain (Model outcomes, Model structure, Treatment efficacy, Evidence considered, Value tool)
- The IVI-RA Technical Expert Panel (TEP) in this survey will score each suggested change according to priority (1-4 with 4 being the highest priority); members must also write in a reason for their choice.

Results Summary:

- Members of the TEP will be provided with anonymized summary tables of the results as well as the reasons that other panelists made the choices they did
- Areas of agreement and disagreement will be identified
- The summary tables will include the following for each question: score provided by each member of the TEP (including the panelist's own rating), mean score, median score, measure of variability, and whether agreement was reached.



Survey 2:

- Panelists retake the survey for questions in which agreement was not reached.
- If they remain outside the consensus, they must write in a reason for this decision.

IVI Prioritization:

- IVI will rank suggested changes (for which there is consensus) by the median/mean 1-4 priority score.

Future updates to the model will be made based on the ranking and financial resources available.

For a detailed description of the model, please refer [here](#).

This survey has 40 questions, and will take approximately 35 minutes to complete. Thank you for your time.

Model assumptions and parameters

Model structure

HAQ rebound

Like most decision-analytic models in rheumatoid arthritis, version 1 of the model measures changes in disease severity (functional status) over time using the Health Assessment Questionnaire (HAQ). HAQ is associated with several model outcomes of interest, including utility, mortality, and costs. One of the most important predictors of cost-effectiveness is the degree to which the HAQ score increases following treatment failure (*See Figure 2; Page 19 of the model description document*). Upon discontinuation of treatment, the HAQ score rebounds by a proportion of the improvement experienced at the end of the initial 6-month period with that treatment. Since no data exists on the size of the HAQ rebound post treatment, we assumed the expected rebound at 85% (with a range of 70% - 100% due to uncertainty) for version 1 of the model. Given the impact of HAQ rebound on model estimates and lack of data, it is an important factor to get consensus on.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
1	The HAQ score should rebound by 100% at treatment discontinuation	1	2	3	4	
2	The current HAQ rebound range from 70% - 100% should be modified to something else.	1	2	3	4	

Progression of DAS28 over time

Some reviewers of the model have suggested to represent the course of disease over time with the disease activity score (DAS) rather than with the HAQ.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
3	Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ.	1	2	3	4	

Radiographic progression

For each targeted DMARD (tDMARD) evaluated in the model, treatment is separated into two distinct phases: an initial phase of up to 6 months, consistent with data reported from randomized controlled trials, and a maintenance phase thereafter until discontinuation. During the initial treatment phase, HAQ is modeled as a change from baseline according to different structural modeling pathways:

H1: Treatment → ACR → HAQ

H2: Treatment → ACR → EULAR → HAQ

H3: Treatment → HAQ

(See Section 7.1; Page 19 of the model description document)

During the public comment period, a reviewer suggested to incorporate the modified Total Sharp Score (mTSS) as a measure of radiographic progression to increase the clinical relevance of the model.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
4	The impact of treatment on HAQ should be modeled through the mTSS (treatment → mTSS → HAQ; H4) as an additional structural pathway that can be selected for model analysis.	1	2	3	4	

HAQ and mortality

In version 1 of the model, the probability of death is simulated as a function of age/sex specific mortality rates from U.S. lifetables, baseline HAQ (i.e, their functional status when they begin taking tDMARDs), and changes in HAQ from baseline. Treatment, therefore, has an indirect effect on mortality through its effect on HAQ. (See Section 8.10; Page 39 of the model description document)

One of the comments received was that although a patient's HAQ score at baseline influences mortality, changes in HAQ over time do not.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
5	Changes in HAQ over time do impact mortality.	1	2	3	4	

Additional utility functions

Individual HAQ scores at a particular point in time were used to simulate EuroQol five dimensions questionnaire (EQ-5D) utility scores (0-1 range), which, in turn, were used to simulate quality-adjusted life-years (QALYs). Version 1 of the model contains two different possible mapping algorithms. 1) The Hernández-Alava et al. (2013) mixture model; and 2) a linear regression analysis by Wailoo et al. (*See Section 7.5, Section 8.9, and Appendices F.1 and F.2; Pages 22, 39, page 39 of the model description document*).

Other (trial-based) algorithms have been reported in the literature (e.g. Bansback et al. 2005, Hurst et al. 2006, Malottki et al. 2011, Brennan et al. 2004, Ducournau et al. 2009, and Boggs et al. 2002) and were suggested by reviewers to incorporate in the model as an alternative modeling option as well.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
6	Additional utility functions beyond those already incorporated in the model should be included.	1	2	3	4	

Treatment switching

In version 1 of the model, simulated patients switch treatment during the initial 6 months if they have a serious infection or if they experience an insufficient treatment effect. The user can choose whether treatment switching due to insufficient efficacy is based on disease activity level or treatment response.

Treatment switching during the initial treatment phase is modeled using 6 different pathways S1-S6.

- S1: Treatment → ACR → Switch
- S2: Treatment → ACR → Δ DAS28 → DAS28 → Switch
- S3: Treatment → ACR → Δ SDAI → SDAI → Switch
- S4: Treatment → ACR → Δ CDAI → CDAI → Switch
- S5: Treatment → Δ DAS28 → DAS28 → Switch
- S6: Treatment → ACR → EULAR → Switch

(*See Pages 20-21 of the model description document*).

For the treatment period beyond 6 months with a given tDMARD (i.e. the maintenance phase), patients remain on treatment until treatment discontinuation or death. Time to treatment discontinuation is based on parametric survival analyses of real-world data.

- When S1 is selected, time to treatment discontinuation is simulated using a single time-to-event curve because we have been unable to obtain curves stratified by ACR response categories.
- When S2-S5 are selected, the time-to-event curves are a function of disease activity level so patients with lower disease activity at the end of the initial treatment phase stay on treatment during the maintenance phase longer, on average.

- When S6 is selected, the time-to-event distributions are stratified by EULAR response category and patients with good response at the end of the initial treatment phase tend to stay on maintenance treatment longer than patients with a moderate response. (*See Section 8.6; Pages 33-37 of the model description document*).

Given the structure of the model, disease severity (i.e. HAQ level) or disease activity level during the maintenance phase does not have a direct effect on discontinuation rates; discontinuation rates are modeled based on observational evidence conditional upon the response / disease activity level at the end of induction treatment (6 months). The question is whether a simulated patient's individual HAQ score at a specific point in time during the maintenance phase of a treatment should determine the probability of discontinuation with that treatment directly.

When defining the treatment sequence, the current version of the model only allows selecting each treatment once. As such, patients cannot continue or restart with the same tDMARD as they experienced a serious infection with.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
7	As an alternative to modeling time-to-treatment-discontinuation during the maintenance phase with a specific tDMARD (>6 months) conditional upon the level of response or disease activity at 6 months with the same tDMARD, the model should provide the option to determine treatment discontinuation conditional upon HAQ or disease activity level during the maintenance phase.	1	2	3	4	
8	The model should provide the option to continue with the same tDMARD despite experiencing a serious infection.	1	2	3	4	

Additional outcomes

The model simulates the health outcomes, costs, and risks associated with treatment. (*See Section 7.8; Page 25 of the model description document*). Depending on the model structure, model outcomes include the following:

- Clinical outcomes during initial treatment phase: ACR response, EULAR response, DAS28, SDAI, CDAI
- Long-term clinical outcomes: HAQ, QALYs
- Adverse events: number of serious infections
- Health care sector costs: drug acquisition and administration costs, general management and monitoring costs, adverse event costs, hospitalization costs
- Non-health care sector costs: productivity losses

Reviewers have suggested that some key outcomes of importance are missing from the model.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
9	Cardiovascular events should be included in the model.	1	2	3	4	
10	Fractures should be included in the model.	1	2	3	4	
11	Adverse events need to be expanded beyond the serious infections currently included in the model.	1	2	3	4	
12	The risk of (reactivation of) a tuberculosis infection should be included in the model.	1	2	3	4	
13	In the model pain scores are estimated based on HAQ scores given their association. These simulated pain scores need to be presented separately as they are an important outcome.	1	2	3	4	
14	Fatigue should be added as an outcome in the model.	1	2	3	4	
15	The burden of care for informal care givers of patients should be incorporated in the model.	1	2	3	4	
16	Risk of long-term nursing home care should be included in the model.	1	2	3	4	

Treatment effects

Additional interventions

The model is capable of simulating a treatment sequence of any arbitrary length. Treatments that can be included in a sequence include conventional DMARDs (cDMARDs), biologic DMARDs, and Janus kinase/STAT (JAK/STAT) pathway inhibitors. The biologic DMARDs and JAK/STAT inhibitors, which we refer to collectively as tDMARDs, included in the current version of the model are:

- Tumor necrosis factor (TNF) inhibitors: etanercept, adalimumab, infliximab, certolizumab, golimumab
- non-TNF inhibitors: abatecept, tocilizumab, rituximab
- Janus kinase/signal transducers and activators of transcription (JAK/STAT) inhibitors: tofacitinib

(See Section 7.8; Page 25 of the model description document).

Several reviewers suggested additional treatment options needed to be included in the model.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
17	The model should provide an option to select biosimilars in addition to the currently available biologic DMARDs.	1	2	3	4	
18	Triple therapy should be included as a treatment option in the model.	1	2	3	4	
19	Sarilumab should be included in the model.	1	2	3	4	
20	Available IV and SC formulations for all biologic DMARDs need to be included in the model, even if there is little or no efficacy data from clinical trials.	1	3	4	4	

HAQ with cDMARD and tDMARD maintenance treatment

With the current version of the model, the natural course of HAQ progression over time in the absence of tDMARDs can be assumed to develop over time according to 1) a constant linear rate; or 2) a non-linear rate (i.e. the rate of HAQ progression declines over time; *See Figure 6 of the model description document*). This second option is currently preferred by IVI.

The trajectory of HAQ over time with tDMARDs develops in a linear fashion, determined by the available evidence available to IVI at time of model development. Since, the model estimates outcomes for a lifetime horizon, the treatment effects in terms of HAQ over time estimated based on the available evidence need to be extrapolated over time. (*See Section 8.3-8.5; Pages 28-33 of the model description document*).

Some commentators have suggested that it is inappropriate to assume that the rate of progression remains constant over time beyond the available data, and that a zero progression rate should be used.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
21	The currently <u>used data</u> for modeling HAQ progression over time in the <u>absence</u> of tDMARDs is appropriate.	1	2	3	4	
22	The current approach to <u>extrapolating</u> HAQ progression beyond the follow-up period of the available evidence in the <u>absence</u> of tDMARDs is appropriate.	1	2	3	4	
23	The currently <u>used data</u> for modeling HAQ progression over time <u>with</u> tDMARDs is appropriate.	1	2	3	4	

24	The current approach to <u>extrapolating</u> HAQ progression beyond the follow-up period of the available evidence <u>with</u> tDMARDs, which assumes linear progression, is appropriate.	1	2	3	4	
25	The rate of HAQ progression should gradually decline toward 0 beyond follow-up periods of the available evidence.	1	2	3	4	

Effect-modifiers

The effect of treatment on ACR response, DAS28, and HAQ at 6 months for tDMARD *naïve* patients are based on a network meta-analyses (NMA) of published randomized controlled trials. Treatment effects for tDMARD *experienced* patients are calculated by multiplying treatment effects for tDMARD naïve patients by a constant value that is the same for all treatments. One limitation of this approach is that we are unable to model the relationship between line of therapy and treatment response; that is, treatment response for a patient who has failed at least one tDMARD is assumed to be reduced by the same amount regardless of line of therapy.

In the current version of the model treatment effects do not vary according to patient characteristics beyond tDMARD naïve/experienced . (Nonetheless, in our R package, treatment effects for each simulated patient can be modeled as a function of any variables chosen by the user.)

(See Section 8.1; Page 26 of the model description document).

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
26	Treatment effects among tDMARD experienced patients should be modeled as a function of the <u>number</u> of previous tDMARDs received	1	2	3	4	
27	Treatment effects among tDMARD experienced patients should be modeled as a function of the <u>type</u> of previous tDMARDs received	1	2	3	4	
28	Treatment effects should be incorporated in the model as a function of patient characteristics.	1	2	3	4	

Duration of maintenance treatment / time-to-treatment discontinuation

In the current version of the model, time to treatment discontinuation in the maintenance phase depends on the pathway (S1-S6) used to model treatment switching and not on type of tDMARD. If S1 is selected, a single treatment discontinuation curve based on an analysis from the CORRONA database is used for all patients. In S2-S5, time to treatment discontinuation is stratified by the level of disease activity, and in S6 treatment duration depends on EULAR response. (See Section 8.6; Page 33-37 of the model description document).

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
29	Additional data sources (beyond CORRONA) for long-term treatment duration should be available in the model for the end-user to choose from.	1	2	3	4	
30	Long-term discontinuation rates should be drug specific.	1	2	3	4	

Drug specific adverse events

In the current version of the model, the rate of a serious infection is assumed to be equal across tDMARDs. (See Section 8.8; Page 38 of the model description document).

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
31	tDMARD specific serious infection rates need to be incorporated.	1	2	3	4	

Incorporating recent real-world/registry data

The current model is based on findings reported in the academic literature. Reviewers have suggested that the model would be improved from a relevance perspective if IVI were able to analyze recent patient-level real-world data for key model parameters.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
32	IVI should spend resources to acquire patient-level real-world data.	1	2	3	4	

Drug prices

It is often argued that the wholesale acquisition cost (WAC) is inappropriate because it does not represent actual transaction prices or include prompt pay or other discounts, rebates, or reductions in price. IVI's current approach to this is to use the WAC price and include the expected discount from the WAC price. That said, some commentators have suggested that the National Average Drug Acquisition Cost (NADAC) is a more appropriate measure of drug price.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
33	The NADAC should be used instead of the WAC to estimate drug acquisition costs.	1	2	3	4	

Other

Facilitate cost-effectiveness analysis of a specific drug

The model simulates the costs and outcomes associated with different treatment sequences for rheumatoid arthritis. As such the model allows evaluating the cost-effectiveness of competing sequential treatment strategies.

If the interest is in a comparison of the cost-effectiveness of particular tDMARDs rather than a comparison of sequences, the subsequent treatments used upon discontinuation of the tDMARD of interest still need to be defined given the lifetime horizon of the analysis. However, one may like to have one "average subsequent treatment sequence" applicable to all tDMARDs to be compared. This provides a natural sequence of subsequent treatments that can help facilitate estimates of differences in lifetime outcomes and costs between the competing tDMARDs, while ensuring that results are not confounded by differences in subsequent treatment sequences across tDMARDs.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
34	The model should allow the end-user to select an "average subsequent treatment sequence" following discontinuation of the initial tDMARD of interest to facilitate tDMARD specific cost-effectiveness analyses.	1	2	3	4	

Present tornado diagrams or other information regarding the contribution of different model components/parameters on estimates of cost-effectiveness

The current version of the model propagates uncertainty in model input parameters and quantifies the uncertainty in model outcomes and cost-effectiveness given the evidence available (i.e. probabilistic sensitivity analysis). However, the model does not allow quantifying the impact of the uncertainty of individual input parameters on the model outcomes and cost-effectiveness.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
35	In addition to probabilistic sensitivity analysis available with the current version of the model, the model should also present tornado diagrams, or another way of presenting the contribution of different model components/parameters on the uncertainty in model outcomes and cost-effectiveness estimates.	1	2	3	4	

Priority of modifications to the model

Of the 35 items presented so far, which 5 do you consider the most important to be implemented in the next version of the model? Please check the corresponding boxes

<input type="checkbox"/> 1	<input type="checkbox"/> 19
<input type="checkbox"/> 2	<input type="checkbox"/> 20
<input type="checkbox"/> 3	<input type="checkbox"/> 21
<input type="checkbox"/> 4	<input type="checkbox"/> 22
<input type="checkbox"/> 5	<input type="checkbox"/> 23
<input type="checkbox"/> 6	<input type="checkbox"/> 24
<input type="checkbox"/> 7	<input type="checkbox"/> 25
<input type="checkbox"/> 8	<input type="checkbox"/> 26
<input type="checkbox"/> 9	<input type="checkbox"/> 27
<input type="checkbox"/> 10	<input type="checkbox"/> 28
<input type="checkbox"/> 11	<input type="checkbox"/> 29
<input type="checkbox"/> 12	<input type="checkbox"/> 30
<input type="checkbox"/> 13	<input type="checkbox"/> 31
<input type="checkbox"/> 14	<input type="checkbox"/> 32
<input type="checkbox"/> 15	<input type="checkbox"/> 33
<input type="checkbox"/> 16	<input type="checkbox"/> 34
<input type="checkbox"/> 17	<input type="checkbox"/> 35
<input type="checkbox"/> 18	

Important determinants of the value of a treatment currently not considered in the model

What components contributing to the value of treatment of rheumatoid arthritis patients are missing in the current version of the model and should IVI consider implementing in the model?

Response:

Value tool and Model Interface

Value Tool: <https://innovationandvalueinitiative.shinyapps.io/ivi-ra/>

Model Interface: <https://innovationandvalueinitiative.shinyapps.io/ivi-ra-expert/>

Target-audience

Reviewers indicated that the target audience of the Value Tool, and Model Interface were not clear.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
36	The Value Tool and Model Interface need a better description of its purpose and target audience	1	2	3	4	

Multi-criteria decision analysis

In IVI's web interfaces, the MCDA is currently based on the following criteria: (i) QALYs, (ii) total health care sector costs, (iii) productivity losses, (iv) number of serious infections, (v) route of administration (oral/injection/infusion) and (vi) time the medication has been on the market.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
37	Rather than presenting the QALYs as one of the dimensions in the MCDA, its components (life expectancy, summary of HAQ over time, adverse event rates) need to be presented in a disaggregated format and allow the end-user to give relative weights to each of these factors independently.	1	2	3	4	
38	In addition to the summary of HAQ over time, ACR response, DAS scores, and pain need to be presented in the MCDA as well.	1	2	3	4	
39	Rather than presenting healthcare sector costs as one of the dimensions in the MCDA, its components (e.g. drug acquisition costs, general management costs, costs related to hospitalizations, and costs due to adverse events) need to be presented in a disaggregated format and allow the end-user to give relative weights to each of these factors.	1	2	3	4	
40	The depth of indications for each tDMARD need to be added as one of the factors in the MCDA.	1	2	3	4	



Delphi Survey #2

Introduction

The IVI-RA model is a part of IVI's Open Source Value Project (OSVP), which is building an open, collaborative, and consensus-based process for the development of tools for value assessment. Models developed by the OSVP process are iterative, evolving as the science of value assessment advances, and as new evidence becomes available. OSVP models are released and updated using a four-step process:

- (1) Public release of the model.
- (2) Invite feedback and suggested improvements to the model in a public comment period.
- (3) Through a 2-part modified Delphi approach, a panel of experts determines which of the evidence-based suggestions for improvement suggested in Step 2 should be implemented by means of peer-review and a formal voting process.
- (4) Revise the model based on the feedback from the technical expert panel in Step 3.

As part of step 3, over the past few weeks, you engaged in Part 1 of a 2-part modified Delphi approach by completing a survey in which you ranked suggested changes to incorporate into the next iteration of the model from the public comments we received. These comments were categorized by domain (model outcomes, model structure, treatment efficacy, evidence considered, value tool).

After synthesizing all responses received, we were able to identify areas of unanimous agreement, areas of general consensus, and areas where agreement was not reached.

This survey serves as the final part of the 2-part modified Delphi approach. The primary goals of this survey are to summarize and inform respondents on areas of agreement from Survey #1, and solicit further feedback for areas where agreement was not reached.

For a detailed description of the model, please refer [here](#).

This survey features 14 questions, and will take approximately 10-15 minutes to complete. Thank you very much for your time.

Areas of unanimous agreement

Based off of responses received in Survey #1, the following were identified as suggested changes that were consistently ranked highly among all respondents, and have been established as top priorities for inclusion in the next version of the RA-Model.

- (1) The option to select biosimilars in addition to the currently available biologic DMARDs. Average score: 4.8/5.0
- (2) The inclusion of triple therapy as a treatment option. Average score: 4.8/5.0
- (3) As an alternative to modeling time-to-treatment-discontinuation during the maintenance phase with a specific tDMARD (>6 months) conditional upon the level of response or disease activity at 6 months with the same tDMARD, providing the option to determine treatment discontinuation conditional upon HAQ or disease activity level during the maintenance phase. Average score: 4.4/5.0
- (4) In addition to probabilistic sensitivity analysis available with the current version of the model, presenting tornado diagrams, or another way of presenting the contribution of different model components/parameters on the uncertainty in model outcomes and cost-effectiveness estimates. Average score: 4.4/5.0

(5) Including additional utility functions beyond those already incorporated in the model. Average score: 4.0/5.0

(6) Including additional data sources (beyond CORRONA) for long-term treatment duration for the end-user to choose from. Average score: 4.0/5.0

1. Please elaborate on any additional comments/rationale you may have regarding any of the 6 suggestions listed above.

Areas of general consensus

The following set of questions were identified as suggestions that were ranked relatively highly as priorities overall, but varied slightly among respondents.

After taking into account the comments that were received regarding each suggestion from Survey #1, please select the answer choice you most closely align with.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
2	<p>Sarilumab should be included in the model. Average score: 4.2/5.0</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> • Yes, any available treatment should be included • Would be based on only RCT • If it is FDA approved 	1	2	3	4	
3	<p>IVI should spend resources to acquire patient-level real-world data. Average score: 4.2/5.0.</p> <p>Comments received regarding the question from Survey #1:</p> <ul style="list-style-type: none"> • These are some areas that could benefit from patient reported outcomes (PROs). In particular, measures of fatigue and measure of HAQ changes before, during, and after treatment switching. • CORRONA would be an excellent source. 	1	2	3	4	

	<ul style="list-style-type: none"> • Always. When possible, do it. • Models need to be updated with real-world data, but I thought this aims to be a tool that allows multiple parties to do this. 					
4	<p>The model should continue to allow changes in HAQ over time to affect mortality. Average score: 4.2/5.0.</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> • I am a bit surprised that changes in HAQ over time don't factor into mortality in the model already. Seems like a good idea. • The HAQ represents the patient's perspective; changes in the HAQ may capture the patient's demand to try a new treatment or remain with the current treatment despite radiologic findings. • Would be great • Not sure what would be gained by this complexity 	1	2	3	4	
5	<p>The model should provide the option to continue with the same tDMARD despite experiencing a serious infection. Average score: 4.0/5.0</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> • I don't think there are guidelines on whether to discontinue or not, and have heard that some people do not switch following hospitalization. Some may have no better alternative. • Seems like this would happen in the real world. • This does occur in the real world. Most people don't but it does happen. Include as an option. 	1	2	3	4	
6	<p>Long-term discontinuation rates should be drug specific. Average score: 4.0/5.0.</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> • If discontinuation rates vary by medication, that would be very helpful information to include. There is a lot 	1	2	3	4	

	<p>of value to avoiding switching when possible.</p> <ul style="list-style-type: none"> Do you have adequate data outside of randomized clinical trials to support this? Make it mechanism of action specific. Model should have this flexibility. 					
7	<p>Treatment effects among tDMARD experienced patients should be modeled as a function of the type of previous tDMARDs received. Average score: 3.8/5.0.</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> If there is sufficient evidence to make this change in the model, it might be useful, though I would have the concern regarding the medication looking less effective (and therefore limiting patient access) 	1	2	3	4	
8	<p>Treatment effects should be incorporated in the model as a function of patient characteristics. Average score: 3.6/5.0</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> If there is sufficient evidence to make this change in the model, it might be useful, though I would have a concern regarding the medication looking less effective (and therefore limiting patient access) 	1	2	3	4	
9	<p>The NADAC should be used instead of the WAC (along with adjustments for rebates and discounts) to estimate drug acquisition costs. Average score: 3.6/4.0</p>	1	2	3	4	

Areas where agreement was not reached

The following set of questions were identified as suggestions where responses consistently varied, and agreement was not reached.

After taking into account the comments that were received regarding each suggestion from Survey #1, please select the answer choice you most closely align with.

		Totally disagree	Somewhat disagree	Somewhat agree	Totally agree	Comment/Rationale
10	<p>Risk of long-term nursing home care should be included in the model. Average score: 4.0/5.0</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> Given the potentially high costs involved with long-term care, this seems like it could be useful to include, if possible. Again, falls are an important determinant of the need for long term care. Only if you have a good paper to support it. Small percentage of patients. I would assume the HAQ would pick this up sufficiently. 	1	2	3	4	
11	<p>Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ. Average score: 3.8/5.0.</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> DAS-28 does not adequately reflect the patient perspective. And it doesn't incorporate issues beyond certain joints. Many patients experience problems in their feet, for example, which is not included in the DAS-28. How sensitive would the model be to this type of change? Is DAS less likely to result in a finding of disease activity than HAQ? Some users may have information on both the DAS-28 and the HAQ and they may wish to explore alternatives from either perspective. 	1	2	3	4	

	<ul style="list-style-type: none"> DAS-28 has more objective markers DAS-28 is a better measure but little data What is the evidence here? What's the correlation between the two? 					
12	<p>Fractures should be included in the model. Average score: 3.8/5.0</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> It seems to me like you would need to also include joint replacement in this category – joints fail and need replacement from RA, but are not always 'fractured' Falls and fractures due to mobility problems are a serious problem regarding both costs and mortality Not likely possible to include well based on data → need to use DAS28 likely to be able to do this; function and fraction are highly correlated. Likely already reflected in QoL measures. Don't have time to event data. Possible to include on event, correlation on HAQ, and effect on QoL. What is the evidence on this? 	1	2	3	4	
13	<p>Treatment effects among tDMARD experienced patients should be modeled as a function of the number of previous tDMARDs received. Average score: 3.6/4.0.</p> <p>Comments received regarding this question from Survey #1:</p> <ul style="list-style-type: none"> I believe there is evidence that medications are less effective as switching continues. If that is true, it seems like this modification to the model would make the results more realistic. However, I'm concerned that making the medication look less effective would result in patients being denied access to medications because the modeled results make it look like they are not as effective. It is intuitively plausible that multiple tDMARD failures would suggest a very stubborn response. 	1	2	3	4	

	<ul style="list-style-type: none"> Used to be useful. Less so now. Number of tDMARDs often a proxy for how well they are doing, rather than being casual. I would imagine that effectiveness declines somewhat over time. 					
--	---	--	--	--	--	--

Please provide any feedback you may have regarding ways to improve the Technical Expert process overall.

Appendix D. Delphi Survey Results

Table 1. Results of Delphi Survey #1: Top Areas of Disagreement Based off Standard Deviation (SD)

<i>IVI RA-Model Technical Expert Panel Delphi Survey #1 Results</i>		
Top areas of disagreement based off of Standard Deviation (SD)	SD	Rank
In the model pain scores are estimated based on HAQ scores given their association. These simulated pain scores need to be presented separately as they are an important outcome.	1.3	1
Cardiovascular events should be included in the model.	1.2	2
tDMARD specific serious infection rates need to be incorporated.	1.2	2
Fatigue should be added as an outcome in the model.	1.1	4
The HAQ score should rebound by 100% at treatment discontinuation.	1.1	5
The risk of (reactivation of) a tuberculosis infection should be included in the model.	0.9	6
Fractures should be included in the model.	0.8	7
Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ.	0.8	7
The rate of HAQ progression should gradually decline toward 0 beyond follow-up periods of the available evidence.	0.8	9
Risk of long-term nursing home care should be included in the model.	0.7	10
The current HAQ rebound range from 70%-100% should be modified to something else.	0.6	11
The impact of treatment on HAQ should be modeled through the mTSS (treatment → mTSS → HAQ; H4) as an additional structural pathway that can be selected for model analysis.	0.5	12
Sarilumab should be included in the model.	0.5	12
The current approach to extrapolating HAQ progression beyond the follow-up period of the available evidence in the absence of tDMARDs is appropriate.	0.5	12
The current approach to extrapolating HAQ progression beyond the follow-up period of the available evidence with tDMARDs, which assumes linear progression, is appropriate.	0.5	12
Treatment effects should be incorporated in the model as a function of patient characteristics.	0.5	12
IVI should spend resources to acquire patient-level real-world data.	0.5	12
The NADAC should be used instead of the WAC to estimate drug acquisition costs.	0.5	12
The model should allow the end-user to select an “average subsequent treatment sequence” following discontinuation of the initial tDMARD of interest to facilitate tDMARD specific cost-effectiveness analyses.	0.5	19
Additional data sources (beyond CORRONA) for long-term treatment duration should be available in the model for the end-user to choose from.	0.5	20

Treatment effects among tDMARD experienced patients should be modeled as a function of the type of previous tDMARDs received.	0.5	20
Available IV and SC formulations for all biologic DMARDs need to be included in the model, even if there is little or no efficacy data from clinical trials.	0.5	20
Additional utility functions beyond those already incorporated in the model should be included.	0.5	20
As an alternative to modeling time-to-treatment-discontinuation during the maintenance phase with a specific tDMARD (>6 months) conditional upon the level of response or disease activity at 6 months with the same tDMARD, the model should provide the option to determine treatment discontinuation conditional upon HAQ or disease activity level during the maintenance phase.	0.4	24
The model should provide the option to continue with the same tDMARD despite experiencing a serious infection.	0.4	24
Adverse events need to be expanded beyond the serious infections currently included in the model.	0.4	24
The burden of care for informal care givers of patients should be incorporated in the model.	0.4	24
The currently used data for modeling HAQ progression over time in the absence of tDMARDs is appropriate.	0.4	24
Long-term discontinuation rates should be drug specific.	0.4	24
In addition to probabilistic sensitivity analysis available with the current version of the model, the model should also present tornado diagrams, or another way of presenting the contribution of different model components/parameters on the uncertainty in model outcomes and cost-effectiveness estimates.	0.4	24
It is important for the model to estimate how changes in HAQ over time affect mortality.	0.4	31
The model should provide an option to select biosimilars in addition to the currently available biologic DMARDs.	0.4	31
Triple therapy should be included as a treatment option in the model.	0.4	31
The currently used data for modeling HAQ progression over time with tDMARDs is appropriate.	0.4	31
Treatment effects among tDMARD experienced patients should be modeled as a function of the number of previous tDMARDs received.	0.4	31

Table 2. Results of Delphi Survey #1: Top Priorities Based Off Average Score

<i>IVI RA-Model Technical Expert Panel Delphi Survey #1 Results</i>		
Top priorities based off of Average Score	Avg. score	Rank
The model should provide an option to select biosimilars in addition to the currently available biologic DMARDs.	3.8	1
Triple therapy should be included as a treatment option in the model.	3.8	1

As an alternative to modeling time-to-treatment-discontinuation during the maintenance phase with a specific tDMARD (>6 months) conditional upon the level of response or disease activity at 6 months with the same tDMARD, the model should provide the option to determine treatment discontinuation conditional upon HAQ or disease activity level during the maintenance phase.	3.8	3
In addition to probabilistic sensitivity analysis available with the current version of the model, the model should also present tornado diagrams, or another way of presenting the contribution of different model components/parameters on the uncertainty in model outcomes and cost-effectiveness estimates.	3.8	3
Additional utility functions beyond those already incorporated in the model should be included.	3.7	5
Additional data sources (beyond CORRONA) for long-term treatment duration should be available in the model for the end-user to choose from.	3.7	5
Sarilumab should be included in the model.	3.5	7
The current approach to extrapolating HAQ progression beyond the follow-up period of the available evidence in the absence of tDMARDs is appropriate.	3.5	7
The current approach to extrapolating HAQ progression beyond the follow-up period of the available evidence with tDMARDs, which assumes linear progression, is appropriate.	3.5	7
Treatment effects should be incorporated in the model as a function of patient characteristics.	3.5	7
IVI should spend resources to acquire patient-level real-world data.	3.5	7
The NADAC should be used instead of the WAC to estimate drug acquisition costs.	3.5	7
The model should allow the end-user to select an “average subsequent treatment sequence” following discontinuation of the initial tDMARD of interest to facilitate tDMARD specific cost-effectiveness analyses.	3.4	13
Treatment effects among tDMARD experienced patients should be modeled as a function of the type of previous tDMARDs received.	3.3	14
Long-term discontinuation rates should be drug specific.	3.3	15
The currently used data for modeling HAQ progression over time in the absence of tDMARDs is appropriate.	3.3	15
The burden of care for informal care givers of patients should be incorporated in the model.	3.3	15
Fractures should be included in the model.	3.3	15
The model should provide the option to continue with the same tDMARD despite experiencing a serious infection.	3.3	15
Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ.	3.3	15
It is important for the model to estimate how changes in HAQ over time affect mortality.	3.2	21
Risk of long-term nursing home care should be included in the model.	3.2	21
The currently used data for modeling HAQ progression over time with tDMARDs is appropriate.	3.2	21

Treatment effects among tDMARD experienced patients should be modeled as a function of the number of previous tDMARDs received.	2.8	24
Adverse events need to be expanded beyond the serious infections currently included in the model.	2.8	25
In the model pain scores are estimated based on HAQ scores given their association. These simulated pain scores need to be presented separately as they are an important outcome.	2.8	25
Cardiovascular events should be included in the model.	2.7	27
tDMARD specific serious infection rates need to be incorporated.	2.7	27
The impact of treatment on HAQ should be modeled through the mTSS (treatment → mTSS → HAQ; H4) as an additional structural pathway that can be selected for model analysis.	2.5	29
Fatigue should be added as an outcome in the model.	2.5	29
The risk of (re)activation of a tuberculosis infection should be included in the model.	2.3	31
Available IV and SC formulations for all biologic DMARDs need to be included in the model, even if there is little or no efficacy data from clinical trials.	2.3	31
The HAQ score should rebound by 100% at treatment discontinuation.	2.3	33
The current HAQ rebound range from 70%-100% should be modified to something else.	2	34
The rate of HAQ progression should gradually decline toward 0 beyond follow-up periods of the available evidence.	2	34

Table 3. Results of Delphi Survey #2: Top Areas of Disagreement Based off Standard Deviation (SD)

<i>IVI-RA Model Technical Expert Panel Delphi Survey #2 Results</i>		
Top areas of disagreement based off of Standard Deviation (SD)	SD	Rank
Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ.	1.1	1
Fractures should be included in the model.	0.7	2
The model should provide the option to continue with the same tDMARD despite experiencing a serious infection.	0.7	2
Treatment effects among tDMARD experienced patients should be modeled as a function of the number of previous tDMARDs received.	0.7	2
The NADAC should be used instead of the WAC (along with adjustments for rebates and discounts) to estimate drug acquisition costs.	0.5	5
Treatment effects should be incorporated in the model as a function of patient characteristics.	0.4	6
The model should continue to allow changes in HAQ over time to affect mortality.	0.4	6
IVI should spend resources to acquire patient-level real-world data.	0.4	6
Sarilumab should be included in the model.	0.4	6

Long-term discontinuation rates should be drug specific.	0.4	6
Treatment effects among tDMARD experienced patients should be modeled as a function of the type of previous tDMARDs received.	0	11
Risk of long-term nursing home care should be included in the model.	0	11

Table 4. Results of Delphi Survey #2: Top Priorities Based Off Average Score

<i>IVI-RA Model Technical Expert Panel Delphi Survey #2 Results</i>		
Top priorities based off of Average Score	Avg. score	Rank
Sarilumab should be included in the model.	3.8	1
Long-term discontinuation rates should be drug specific.	3.8	1
The model should continue to allow changes in HAQ over time to affect mortality.	3.8	1
The NADAC should be used instead of the WAC (along with adjustments for rebates and discounts) to estimate drug acquisition costs.	3.3	4
IVI should spend resources to acquire patient-level real-world data.	3.3	4
The model should provide the option to continue with the same tDMARD despite experiencing a serious infection.	3.2	6
Treatment effects among tDMARD experienced patients should be modeled as a function of the type of previous tDMARDs received.	3	7
Risk of long-term nursing home care should be included in the model.	3	7
Treatment effects among tDMARD experienced patients should be modeled as a function of the number of previous tDMARDs received.	3	7
Fractures should be included in the model.	2.8	10
Progression of DAS-28 over time should be included in the model as an alternative to progression of disease according to HAQ.	2.8	10
Treatment effects should be incorporated in the model as a function of patient characteristics.	2.8	10

References

1. Incerti, D.J., Jeroen, *A Description of the IVI-RA Model v1.0.* 2017, Innovation and Value Initiative.
2. Wolfe, F., et al., *Predicting mortality in patients with rheumatoid arthritis.* Arthritis & Rheumatism: Official Journal of the American College of Rheumatology, 2003. **48**(6): p. 1530-1542.